

Certificate of Mailing	
Date of Deposit <u>October 28, 2001</u>	Label Number: <u>EL509218278US</u>
I hereby certify under 37 C.F.R. § 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" with sufficient postage on the date indicated above and is addressed to BOX PATENT APPLICATION, Assistant Commissioner for Patents, Washington, D.C. 20231.	
Guy E. Beardsley	<i>Guy E Beardsley</i>
Printed name of person mailing correspondence	Signature of person mailing correspondence

APPLICATION  
FOR  
UNITED STATES LETTERS PATENT

APPLICANT : Mehran M. Khodadoust

TITLE : COMPOSITIONS AND METHODS FOR THE DISCOVERY  
AND SELECTION OF BIOLOGICAL INFORMATION

5                    COMPOSITIONS AND METHODS FOR THE DISCOVERY  
                      AND SELECTION OF BIOLOGICAL INFORMATION

CROSS-REFERENCE TO RELATED APPLICATION

                      This application is a continuation-in-part of U.S. Utility Application No.  
09/908,305, filed July 17, 2001, which is still pending and is a continuation-in-part of  
10 U.S. Utility Application No. 09/697843, filed October 27, 2000.

BACKGROUND OF THE INVENTION

                      The present invention relates to compositions and methods which can be used to  
obtain biological information from cells. Such information, which includes the regulatory  
15 pathways and components utilized by ligands and cytokines to regulate the expression of  
genes in a variety of specific cells and tissue types, can be used to maximize a drug  
targeting strategy for the selection of drug candidates and for library screening. Vectors  
are modified to introduce regulatory or reporter genes into regulated loci within the  
genome of a desired cell line. Such vectors utilize positive and negative selection  
20 markers for the identification and delivery of genes to the regulated loci. Individual  
isolated cell clones that express the reporter marker in a stimulation-dependent manner  
are used as reporter cell lines to test the functional activity of the ligand in the cells. The  
cell clones can be used to identify key regulatory factors, such as promoters and  
enhancers that are dependent on stimulation by the ligand and genes regulated by the  
25 ligand. Genes that can control the response of a regulated loci to stimulation by a ligand  
or cytokine in the selected cell or tissue type can also be identified.

                      Modern drug discovery techniques are increasingly based on genomics and  
depend upon the identification of specific genomic targets and regulatory pathways.  
These genomic targets include the specific genes of interest and their cell-specific  
30 regulatory control elements, such as enhancers and promoters. The expression of  
regulatory factors occurs in a variety of specific cell and tissue types and involves a  
multiplicity of pathways in an organism. An inhibitor or antagonist for a given factor can

have unintended consequences if these complexities are not fully explored and resolved. See, for instance, Khodadoust *et al.*, *Blood*, 92, No. 7, pages 2399-2409 (1998), which describes the distinct regulatory mechanisms for IFN- $\alpha/\beta$  and IFN- $\gamma$ -mediated induction of the Ly-6E gene in B cells. Through deletion analysis, it was found that a cooperative  
 5 interaction exists between physically disparate regulatory regions of the gene. This indicates the complexity involved in achieving cell-type specificity in IFN-mediated gene regulation and is an example of the complexity involved in dealing with these effects.

Regulation of gene expression can be investigated by the integration of promoterless selectable marker genes into the chromosomal loci of cells and the  
 10 subsequent identification of the active loci. This type of "induction trap" strategy has been used to identify specific enhancers, promoters, and other regulatory elements of genes of interest. Induction gene trap vectors, which generate spliced fusion transcripts between the reporter gene and the endogenous gene present at the site of integration, are used to identify regulated gene loci. This approach can be used to distinguish between  
 15 genes involved in specific regulatory pathways and the "housekeeping" genes, which are turned on independently of activation by a ligand. The genes regulated by a ligand would be implicated in regulatory pathways of interest that can be harnessed in drug development.

Gene trap vectors, which generally consist of a splice-acceptor site located  
 20 upstream from a reporter gene, target the introns of the eukaryotic genome. Integration of the reporter into an intron results in a fusion transcript containing mRNA from the endogenous gene and from the reporter gene sequence. The use of an IRES site between the splice acceptor and the reporter gene of a gene trap vector means that the reporter gene product and the endogenous gene product need not be fusion products, thereby  
 25 increasing the likelihood that integration of the vector will result in expression of the reporter gene product. Gene entrapment vectors, or gene trap vectors, are tools which are frequently used for gene discovery and elucidation. These vectors can be employed to identify developmentally regulated genes.

U.S. Patent No. 5,922,601 describes an induction gene trap construct used for the  
 30 identification of genes that are regulated upon the occurrence of a cellular transition event. The construct contains a functional splice acceptor, a translation stop sequence, an

internal ribosome entry site ("IRES"), and a promoterless protein coding sequence encoding a polypeptide providing positive and negative selection traits. The positive and negative selection traits can be introduced by employing nucleic acid encoding a single protein whose expression (or non-expression) can be detected as a positive or negative selection trait. Typical proteins of this type include neomycin phosphotransferase and thymidine kinase. The construct is incorporated in a vector which is introduced into a cell, and the expression of the positive and negative selection traits before and after occurrence of the transition event is detected by means of drug selection. The transition event is typically the transition from an undifferentiated cell to a differentiated cell. The gene trap vector of this reference allows for the selection of genes at the cell populations in which a trapped locus is either active or becomes inactive as a result of a cellular transition event.

Mainguy *et al.*, *Nature Biotechnology*, 18, pages 746-749 (2000) describes vectors for use as induction gene traps to identify homeoprotein target genes. The vectors used in this reference include the PT2 bicistronic gene containing the *lacZ* gene fused to a splice acceptor, a thymidine kinase gene driven by an IRES, and a Neo gene under the control of the phosphoglycerate kinase promoter. The PT2 gene trap vector allows the use of gancyclovir for selecting against integration of the vector into constitutively active genes. Hence, subsequent activation allows for the selection and isolation of regulated genes. Using this vector, an embryonic stem cell gene trap library was constructed and screened for activity towards engrailed homeodomain protein. See, also, European Patent No. 902,092, which discloses a similar procedure.

U.S. Patent No. 5,928,888 describes an induction gene trap method, identifying active genomic polynucleotides for identifying proteins and compounds that modulate genomic polynucleotides. The reference achieves this result by inserting a beta-lactamase polynucleotide into the genome of a eukaryotic cell. The cell is then contacted with a predetermined amount of an agent which activates the beta-lactamase, and the amount of beta-lactamase activity is measured. The expressing and non-expressing cells are separated, and the integration of the beta-lactamase gene in the genome of the cells is determined. The reference states that the beta-lactamase reporter provides a mechanism for preparing a genomic integration assay for drug discovery in a high throughput format.

See, also, Whitney *et al.*, *Nature Biotechnology*, 16, pages 1329-1333 (1998), which describes a genome-wide functional assay for the rapid isolation of cell clones and genetic elements responsive to specific stimuli. This assay uses a promoterless beta-lactamase reporter gene transfected into a human T-cell line to generate a living library of reporter-tagged clones. Flow cytometry and fluorogenic substrates were used to identify patterns of regulation associated with specific genes.

PCT published application, publication number WO 99/02719, discloses methods and DNA constructs which can be used for the detection and manipulation of a target eukaryotic gene whose expression is restricted to specific tissue or specialized cell types.

According to this reference, an embryonic stem cell is transformed with a vector containing a first component under the control of a promoter which has restricted expression in a particular cell or tissue type. The stem cell is also transformed with a gene trap vector encoding a second indicator component. The indicators act in a complementary way to produce a detectable signal, such as the omega and alpha components of  $\beta$ -galactosidase which combine to form the complete enzyme. Measurement of the detectable enzyme indicates that the gene of the gene trap vector has been integrated into the genome of the selected cell type.

The activity of a ligand in an organism involves a multiplicity of regulatory pathways depending on the specific cell or tissue type under investigation. For instance, a particular growth factor, such as stem cell factor ("SCF"), can activate cells unrelated to the cell type under investigation, such as mast cells. This lack of specificity, redundancy, and potential toxicity creates a complication when using these factors as protein therapeutics or when developing, for instance, inhibitors, antagonists, or agonists to these factors, since these factors, inhibitors, or antagonists may act on the unrelated cells in unfavorable ways.

It is therefore an objective of this invention to provide a method for characterizing the response of an organism to a ligand by identifying the genes and regulatory mechanisms, in specific cells and tissues, which are activated by the ligand. It is also an object of this invention to discover genes, cells containing the genes and gene regulatory mechanisms which can be used to screen libraries of compounds to find potential drug candidates of interest.

## SUMMARY OF THE INVENTION

The present invention relates broadly to the identification of cellular pathways utilized by ligands of interest to regulate specific genomic loci, and to polynucleotide segments which can be incorporated into induction gene trap vectors in a manner such that they would be operably linked to the regulated loci in the specific transfected eukaryotic cells. The use of the incorporated polynucleotides, or the proteins coded by the polynucleotides, for the selection and identification of cells which are responsive to one or more stimulatory agents, provides a panel of cell clones which can be used to dissect regulatory mechanisms. These polynucleotides and proteins can be used to screen a library of drug candidates to obtain promising therapeutic agents for further evaluation. In addition, the protein coded by the polynucleotide can be used as a means to influence the expression of a gene trap vector. This allows for the isolation and identification of the genes, which affect the up or down regulation of the genomic loci in such cells. These genes may be used to directly screen libraries for therapeutic agents. The genes themselves may have therapeutic applications in, for example, combination therapies or drug discovery.

The current use of protein therapeutics, such as ligands, cytokines or antibodies, and approaches for selecting antagonists, inhibitory agents, and agonists against these factors for use in drug therapy, fail to take into account ligand redundancies and the selectivity of specific cell and tissue types with respect to regulatory pathways under investigation. Therapeutics selected in this way can have unintended consequences, such as toxicity and the selection of inhibitory agents which have an adverse impact on the regulation of cells and tissue not implicated in the disease. Additional complexity is introduced when it is desired to use more than one ligand in combination therapies to achieve a desired therapeutic effect. The use of multiple ligands can involve numerous regulatory pathways in divergent cell types, and this can also have unintended consequences for the regulation of cells which are implicated in the disease under investigation. However, there are currently no simple solutions to this problem.

The vectors and methods of this invention can be used to generate a panel or library of cells under the control of regulatory elements for one or more stimulatory

agents of interest. This collection of cells can be used, in turn, as targets to screen libraries of drug candidates to identify potential therapeutics having a high level of specificity, safety, and effectiveness. In addition, genes which are under the control of the regulatory elements and are responsive to specific stimulatory agents of interest, and genes which regulate the genomic loci and effect the expression of such loci, can be identified, isolated and characterized. These genes, or their regulatory factors, can also be used to screen libraries of drug candidates in a variety of assay formats, such as cell based assays. Stimulatory agents of interest include insulin, stem cell factor ("SCF"), vascular endothelial growth factor ("VEGF"), IL-2, IL-3, IL-6, IgE, FGF-1, FGF-2, FGF-3, TGF- $\beta$ , TNF- $\beta$ , and TNF- $\alpha$ .

Accordingly, in one aspect, the present invention includes nucleic acid constructs and vectors containing the constructs, for infecting cells and for generating a panel of cell clones which can be used as screening tools to screen libraries of candidate drug molecules.

According to this aspect, in one embodiment a nucleic acid construct for use in preparing a vector for generating a panel of cells comprises the following elements in downstream (5' to 3') sequence: a cassette containing an internal ribosome entry site; a transactivator polypeptide coding sequence encoding a polypeptide, said polypeptide acting as a regulator unit to one or more regulatory elements contained in a genomic loci in a particular cell or cell type of interest, said transactivator polypeptide being responsive to one or more regulatory elements contained in a genomic loci in a cell of interest; a translation stop sequence; an internal ribosome entry site; a reporter element responsive to at least one stimulatory agent; and a translation stop sequence.

In preferred features of this embodiment of the invention, the reporter element can be an enzyme, such as secreted alkaline phosphatase, Luciferase<sup>TM</sup>, or green fluorescent protein ("GFP"), the marker polypeptide can be a the promoterless protein coding sequence, such as a tetracycline regulator unit (tTA). A nucleic acid cassette containing these elements can be incorporated into an induction gene trap vector containing a splice acceptor site; an internal ribosome entry site; a marker polypeptide coding sequence encoding a polypeptide providing selection traits and being responsive to one or more regulatory elements contained in a genomic loci in a particular cell or cell type of interest;

and a translation stop sequence. The marker polypeptide can be a fusion protein with positive and negative selection traits. Negative selection traits can be provided in situations whereby the expressed gene leads to the elimination of the host cell, frequently in the presence of a nucleoside analog, such as gancyclovir. Positive selection traits can be provided by drug resistance genes. Suitable negative selection markers include, for example, DNA sequences encoding Hp<sub>r</sub>t, g<sub>p</sub>t, HSV-tk, diphtheria toxin, ricin toxin, and cytosine deaminase. Suitable positive selection traits include, for example, DNA sequences encoding neomycin resistance, hygromycin resistance, histidinol resistance, xanthine utilization, Zeocin resistance, and bleomycin resistance. A particularly preferred fusion protein is a fusion protein encoding Tk-Zeo.

This nucleic acid construct can be incorporated into a vector, such as a viral vector, and preferably a retroviral vector, to transfect cells of interest. This can be accomplished by introducing the vector into a medium containing the cells using techniques known to those skilled in the art. Suitable techniques are described in U.S. Patent No. 5,922,601, the disclosure of which is incorporated herein by reference in its entirety. Not all of the cells will be successfully transfected, meaning that the vector will not be integrated into the genomic loci of the cell. Successful integration events can be selected for using a drug selection compound, such as zeocin. If the vector contains a zeocin resistant gene, the zeocin will serve to kill the cells in which the vector has not been successfully integrated into the genome of the cell.

Once cells which have been successfully transfected, and the vector has been operably integrated into the genome of the cell, the cells are selected for activity with respect to specific stimulatory agents. Such activity can include those cells in which regulatory factors, such as enhancers and promoters, have been turned on by the stimulatory agent, and those cells in which the appropriate regulatory factors have been turned off by the stimulatory agent. Each of these cases will involve a different selection protocol.

To select for cells which have been turned on by the stimulatory agent, the stimulatory agent is introduced into the culture medium with zeocin, a positive selective agent, or another appropriate drug selection agent. The stimulatory agent can be added to the medium either prior to, subsequent to, or together with the drug selection agent. The

stimulatory agent activates the promoterless first marker polypeptide coding sequence in the vector, which encodes a polypeptide conferring drug resistance to the drug selection agent. Regulatory factors present in housekeeping genes present in the cell loci can also be activated independently of the stimulatory agent. However, housekeeping genes are not specific for the stimulatory agent and must be eliminated. This is essential in order to obtain isolated cultures of cells specific for the stimulatory agent. This can be accomplished by adding a negative selection agent such as gancyclovir, for example, which is acted on by the TK (thymidine kinase) gene to eliminate cells expressing the protein. Those cells remaining in the cell culture after treatment with gancyclovir are clones with the vector inserted into the genomic loci which are turned on by the particular stimulatory agent. The expression of the reporter (SEAP) gene, which is turned on by the stimulatory agent, can be used as the readout, allowing the cells to be used directly as drug targets to screen libraries of compounds, or treated with other stimulatory agents in the same manner indicated above in order to identify clones which are capable of being turned on by more than one stimulatory agent.

To select for cells which have been turned off by the stimulatory agent, Zeocin is first introduced into the cell culture medium to eliminate those cells which do not have the vector integrated into the genome of the cell. The stimulatory agent and gancyclovir are then added to a medium containing the cells, and the housekeeping genes which are active in the cell are eliminated. However, the cell clones which are turned off upon treatment with the stimulatory agent remain in the culture. These cells can also be used as drug targets, or treated with other stimulatory agents as indicated above for the selection of genes which are turned on by the stimulatory agent. As will be appreciated, other possible combinations for cell selection using several stimulatory agents can be readily envisioned.

In another embodiment, the nucleic acid construct can contain a promoter regulating the expression of the marker polypeptide coding sequence. The promoter acts on one of the polynucleotide sequences which comprise the marker polypeptide encoding region of the construct. Preferably, the polynucleotide sequence encodes a positive selection marker, such as Zeocin. In this embodiment, the promoter is phosphoglycerate kinase ("PGK").

This vector can also be used to generate a panel or library of cells under the control of regulatory elements activated by one or more stimulatory agents of interest using the method set forth below.

To select for cells which have been turned on by the stimulatory agent, the cells  
 5 are transformed with the vector containing a promoter for the selection marker. A selection drug, such as zeocin, is introduced into the culture medium to eliminate cells in which the vector has not been integrated into the genomic loci. The culture medium is changed, and gancyclovir is introduced to eliminate housekeeping genes which are active and express Tk. A stimulatory agent is then added to the medium, and the amount of  
 10 secreted alkaline phosphatase is measured as a positive indicator of the presence of cells which are responsive to the stimulatory agent. Those cells generating SEAP and which are turned on by the stimulatory agent are selected, separated, and used as drug targets.

To select for cells which have been turned off by the stimulatory agent, zeocin is introduced into the cell culture medium to eliminate those cells which do not have the  
 15 vector integrated into the genome of the cell. The stimulatory agent and gancyclovir are then added to a medium containing the cells, and the housekeeping genes which are active in the cell are eliminated. The medium is changed, and the cell clones which produce secreted alkaline phosphatase in the absence of the stimulatory agent are selected by measuring the amount of secreted alkaline phosphatase produced. These cells can also  
 20 be used as drug targets, or treated with other stimulatory agents to prepare cells which are specific for more than one stimulatory agent.

In another aspect, a method is provided for selecting cells and cell clones from a medium containing a collection of cells having a specific response to a selected stimulatory agent. The method involves transforming eukaryotic cells with an induction  
 25 gene trap vector to operably integrate the vector into the genome of the cell. The vector can be any vector, such as the vectors described previously, which includes a marker polypeptide coding sequence and a reporter element. Optionally, the vector can also include a transactivator coding sequence. Cells are selected having one or more regulatory elements which activate the reporter polypeptide and, if present, the  
 30 transactivator. Cells which are specifically activated by the stimulatory agent are

selected. These cells can then be used in cell-based assays for screening libraries of potential therapeutic agents.

In yet another aspect, a method is provided for selecting cells and cell clones from a medium containing a collection of cells having a specific response to a selected stimulatory agent. The method involves transforming eukaryotic cells with an induction gene trap vector to operably integrate the vector into the genome of the cell. The vector can be any vector, such as the vectors described previously, which includes a marker polypeptide coding sequence and a reporter element. Optionally, the vector can also include a transactivator coding sequence. Cells are selected having one or more regulatory elements which activate the reporter gene and, if present, the transactivator. Cells which are specifically inactivated by the stimulatory agent are selected. These cells can then be used in cell-based assays for screening libraries of potential therapeutic agents.

In a further aspect of the invention, a gene trap vector can be used to identify genes that lead to transcriptional control, or which up regulate or down regulate the genomic loci of the isolated cell clones. These vectors can be viral vectors, and preferably retroviral vectors, which are integrated into the genomic loci of the cell.

This vector contains a nucleic acid construct including, in 5' to 3' sequence: a minimal promoter sequence containing a transactivator regulatory element, such as a tetracycline responsive element; a protein coding sequence encoding a marker polypeptide providing positive selection traits, the protein being responsive to the transactivator regulatory elements; an internal ribosome entry site; and a functional splice donor site. The tetracycline regulator unit introduced by the induction gene trap vector generates a protein which is activated or repressed by tetracycline in an "on/off" mode and binds to the tetracycline responsive element within the minimal promoter sequence of the gene trap vector and leads to its transcriptional control.

When the cell is contacted with a stimulatory agent, the tetracycline regulator protein is turned on. This protein binds to the minimal promoter sequence containing the tetracycline responsive element, causing the promoter ("TRE<sub>Pcmv</sub>") to activate and cause transcription of a gene downstream of the promoter. This gene, in turn, up regulates or down regulates the genomic loci, causing the tetracycline regulator unit to express

protein, thereby activating the TRE<sub>P<sub>CMV</sub></sub> promoter/transactivator to transcribe additional copies of the gene, and so on. Eventually, as a result of this feedback process, enough genetic material will be generated to be detected and identified.

5 The net result of this process is to isolate genes that up regulate or down regulate the loci of cells which respond to stimulation by one or more stimulatory agents. These genes can then be used as drug targets or as potential therapeutics (*e.g.*, as gene therapy constructs or antisense molecules). Regulatory elements contained in the gene, such as promoters and enhancers, can also be isolated, characterized, and used for drug discovery.

10 The use of the cells, genes, and regulatory elements of this invention to select drug candidates for use as therapeutic agents is conventional in the art. For instance, the cells can be used in live cell screening assays which are effective to evaluate the specificity, toxicity and dosage of a selected therapeutic agent. If live cell assays are not available, conventional assay screening techniques can be used.

15 In another aspect, the invention provides a method of selecting for one or more cells having a specific response to a stimulatory agent of interest. This method involves inserting a vector including a cassette having a positive selection marker, a negative selection marker, and a reporter gene into eukaryotic cells under conditions that result in the integration of the cassette into the genome of the cells. The reporter gene is operably linked to an endogenous regulatory element in at least one cell. Cells in which expression  
20 of the reporter gene is specifically activated by the stimulatory agent are selected. In particular embodiments, this selection step may involve incubating the cells in the presence of the stimulatory agent and a positive selection agent and incubating the cells under conditions in which a negative selection agent is present and the stimulatory agent is absent. In other embodiments, the selection step involves incubating the cells in the  
25 presence of a positive selection agent, incubating the cells in the presence of a negative selection agent, incubating the cells in the presence of the stimulatory agent, and selecting the cells that express the reporter gene in the presence of the stimulatory agent. In yet other embodiments, the vector does not contain a promoter operably linked to the reporter gene. In various embodiments, the vector further includes a nucleic acid segment  
30 encoding a transactivator polypeptide (*e.g.*, tTA) that is integrated into the genome of the cells. The nucleic acid segment encoding a transactivator polypeptide may be operably

linked to a promoter in the vector or may not be operably linked to a promoter in the vector. Desirably, the nucleic acid encoding the transactivator polypeptide is integrated into the genome of the cells under the control of an endogenous regulatory element. Optionally, the methods may include identifying the regulatory element that is activated by the stimulatory agent. In various embodiments, the positive selection marker is operably linked to a prokaryotic promoter in the cassette that integrates into the genome of the eukaryotic cells. In this case, the regulatory element activated by the stimulatory agent may be identified by (i) inserting a nucleic acid that includes the positive selection marker and a segment of the eukaryotic genome flanking the integrated cassette into bacterial cells under conditions that allow the selection of bacterial cells expressing the positive selection marker under the control of the prokaryotic promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected bacterial cells, and (iii) sequencing the amplified segment. In other embodiments, the positive selection marker is operably linked to a yeast promoter in the cassette that integrates into the genome of the eukaryotic cells. In this case, the regulatory element activated by the stimulatory agent may be identified by (i) inserting a nucleic acid that includes the positive selection marker and a segment of the eukaryotic genome flanking the integrated cassette into yeast cells under conditions that allow the selection of yeast cells expressing the positive selection marker under the control of the yeast promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected yeast cells, and (iii) sequencing the amplified segment.

In a related aspect, the invention provides another method of selecting for one or more cells having a specific response to a stimulatory agent of interest. This method involves inserting a vector including a cassette having a positive selection marker, a negative selection marker, and a nucleic acid segment encoding a transactivator polypeptide into eukaryotic cells under conditions that result in the integration of the cassette into the genome of the cells. The transactivator polypeptide is operably linked to an endogenous regulatory element in at least one cell. Cells in which expression of the transactivator polypeptide is specifically activated by the stimulatory agent are selected. In particular embodiments, this selection step may involve incubating the cells in the presence of the stimulatory agent and a positive selection agent and incubating the cells

under conditions in which a negative selection agent is present and the stimulatory agent is absent. In other embodiments, the selection step involves incubating the cells in the presence of a positive selection agent, incubating the cells in the presence of a negative selection agent, incubating the cells in the presence of the stimulatory agent, and selecting

5 the cells that express the transactivator polypeptide in the presence of the stimulatory agent. In yet other embodiments, the vector does not contain a promoter operably linked to the nucleic acid segment encoding a transactivator polypeptide. In various embodiments, the vector further includes a reporter gene that is integrated into the genome of the cells. The reporter gene may be operably linked to a promoter in the

10 vector or may not be operably linked to a promoter in the vector. Desirably, the reporter gene is integrated into the genome of the cells under the control of an endogenous regulatory element. The methods may optionally include identifying the regulatory element that is activated by the stimulatory agent. In various embodiments, the positive selection marker is operably linked to a prokaryotic promoter in the cassette that

15 integrates into the genome of the eukaryotic cells. In this case, the regulatory element activated by the stimulatory agent may be identified by (i) inserting a nucleic acid that includes the positive selection marker and a segment of the eukaryotic genome flanking the integrated cassette into bacterial cells under conditions that allow the selection of bacterial cells expressing the positive selection marker under the control of the

20 prokaryotic promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected bacterial cells, and (iii) sequencing the amplified segment. In other embodiments, the positive selection marker is operably linked to a yeast promoter in the cassette that integrates into the genome of the eukaryotic cells. In this case, the regulatory element activated by the stimulatory agent may be identified by (i) inserting a

25 nucleic acid that includes the positive selection marker and a segment of the eukaryotic genome flanking the integrated cassette into yeast cells under conditions that allow the selection of yeast cells expressing the positive selection marker under the control of the yeast promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected yeast cells, and (iii) sequencing the amplified segment.

30 In a related aspect, the invention provides yet another method of selecting for one or more cells having a specific response to a stimulatory agent of interest. This method

includes inserting a vector including a cassette having a positive selection marker, a negative selection marker, and a reporter gene into eukaryotic cells under conditions that result in integration of the cassette into the genome of the cells. The reporter gene is operably linked to an endogenous regulatory element in at least one cell. Cells in which

5 expression of the reporter gene is specifically inactivated by the stimulatory agent are selected. In various embodiments, this selection involves incubating the cells in the presence of a positive selection agent and incubating the cells in the presence of the stimulatory agent and a negative selection agent. In yet other embodiments, the vector does not contain a promoter operably linked to the reporter gene. In various

10 embodiments, the vector further includes a nucleic acid segment encoding a transactivator polypeptide (*e.g.*, tTA) that is integrated into the genome of the cells. The nucleic acid segment encoding a transactivator polypeptide may be operably linked to a promoter in the vector or may not be operably linked to a promoter in the vector. Desirably, the nucleic acid encoding the transactivator polypeptide is integrated into the genome of the

15 cells under the control of an endogenous regulatory element. In other embodiments, the method also includes identifying the regulatory element that is inactivated by the stimulatory agent. In various embodiments, the positive selection marker is operably linked to a prokaryotic promoter in the cassette that integrates into the genome of the eukaryotic cells. In this case, the regulatory element inactivated by the stimulatory agent

20 may be identified by (i) inserting a nucleic acid that includes the positive selection marker and a segment of the eukaryotic genome flanking the integrated cassette into bacterial cells under conditions that allow the selection of bacterial cells expressing the positive selection marker under the control of the prokaryotic promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected bacterial cells, and (iii)

25 sequencing the amplified segment. In other embodiments, the positive selection marker is operably linked to a yeast promoter in the cassette that integrates into the genome of the eukaryotic cells. In this case, the regulatory element inactivated by the stimulatory agent may be identified by (i) inserting a nucleic acid that includes the positive selection marker and a segment of the eukaryotic genome flanking the integrated cassette into yeast cells

30 under conditions that allow the selection of yeast cells expressing the positive selection marker under the control of the yeast promoter, (ii) amplifying the segment of the

eukaryotic genome that is inserted into the selected yeast cells, and (iii) sequencing the amplified segment.

In a related aspect, the invention provides still another method of selecting for one or more cells having a specific response to a stimulatory agent of interest. This method includes inserting a vector including a cassette having a positive selection marker, a negative selection marker, and a nucleic acid segment encoding a transactivator polypeptide into eukaryotic cells under conditions that result in integration of the cassette into the genome of the cells. The nucleic acid segment encoding a transactivator polypeptide is operably linked to an endogenous regulatory element in at least one cell.

Cells in which expression of the transactivator polypeptide is specifically inactivated by the stimulatory agent are selected. In various embodiments, this selection involves incubating the cells in the presence of a positive selection agent and incubating the cells in the presence of the stimulatory agent and a negative selection agent. In yet other embodiments, the vector does not contain a promoter operably linked to the nucleic acid segment encoding a transactivator polypeptide. In various embodiments, the vector further includes a reporter gene that is integrated into the genome of the cells. The reporter gene may be operably linked to a promoter in the vector or may not be operably linked to a promoter in the vector. Desirably, the reporter gene is integrated into the genome of the cells under the control of an endogenous regulatory element. The method may optionally include identifying the regulatory element that is inactivated by the stimulatory agent. In various embodiments, the positive selection marker is operably linked to a prokaryotic promoter in the cassette that integrates into the genome of the eukaryotic cells. In this case, the regulatory element inactivated by the stimulatory agent may be identified by (i) inserting a nucleic acid that includes the positive selection marker and a segment of the eukaryotic genome flanking the integrated cassette into bacterial cells under conditions that allow the selection of bacterial cells expressing the positive selection marker under the control of the prokaryotic promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected bacterial cells, and (iii) sequencing the amplified segment. In other embodiments, the positive selection marker is operably linked to a yeast promoter in the cassette that integrates into the genome of the eukaryotic cells. In this case, the regulatory element inactivated by the stimulatory agent

may be identified by (i) inserting a nucleic acid that includes the positive selection marker and a segment of the eukaryotic genome flanking the integrated cassette into yeast cells under conditions that allow the selection of yeast cells expressing the positive selection marker under the control of the yeast promoter, (ii) amplifying the segment of the  
 5 eukaryotic genome that is inserted into the selected yeast cells, and (iii) sequencing the amplified segment.

In another aspect, the invention provides a method for identifying a nucleic acid of interest that encodes a protein that modulates the activity of a regulatory element in a cell. This method includes inserting a first vector including a first cassette having a positive  
 10 selection marker, a negative selection marker, a reporter gene, and a nucleic acid segment encoding a transactivator polypeptide into eukaryotic cells under conditions that result in integration of the first cassette into the genome of the cells. The reporter gene is operably linked to an endogenous regulatory element in at least one cell, or the reporter gene is operably linked to a regulatory element in the first vector. A second vector including a  
 15 second cassette having a promoter operably linked to a responsive element that is responsive to the transactivator polypeptide is also inserted into the cells under conditions that result in integration of the second cassette into the genome of the cells. The promoter is operably linked to an endogenous nucleic acid of interest encoding a protein that modulates (*i.e.*, increases or decreases) the activity of the regulatory element in at least  
 20 one cell. Cells that have an altered level of reporter gene expression under conditions that activate the transactivator polypeptide are selected. Desirably, the nucleic acid of interest from at least one selected cell is identified. The nucleic acid of interest is determined to encode a protein that activates the regulatory element if the cells have increased reporter gene expression under conditions that activate the transactivator polypeptide. Or the  
 25 nucleic acid of interest is determined to encode a protein that inactivates the regulatory element if the cells have decreased reporter gene expression under conditions that activate the transactivator polypeptide. In desirable embodiments, the transactivator polypeptide is tTA, and the responsive element comprises a tetracycline responsive element. In various embodiments, the first vector contains a regulatory element that was identified as  
 30 being regulated by a stimulatory agent of interest. For example, the methods of the invention may be used to identify an endogenous regulatory element that is regulated by a

stimulatory agent, and then this regulatory element may be cloned into the first vector using standard methods and used to identify endogenous nucleic acids encoding proteins that modulate the regulatory element. In other embodiments, the second vector further includes a positive selection marker which is integrated into the genome of the cells. In still other embodiments, the positive selection marker in the second vector (denoted the second positive selection marker) is operably linked to a prokaryotic promoter in the second cassette that integrates into the genome of the eukaryotic cells. In this case, the nucleic acid encoding a protein that modulates the activity of the regulatory element may be identified by (i) inserting a nucleic acid that includes the second positive selection marker and a segment of the eukaryotic genome flanking the second integrated cassette into bacterial cells under conditions that allow the selection of bacterial cells expressing the second positive selection marker under the control of the prokaryotic promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected bacterial cells, and (iii) sequencing the amplified segment. In yet other embodiments, the positive selection marker in the second vector is operably linked to a yeast promoter in the second cassette that integrates into the genome of the eukaryotic cells. In this case, the nucleic acid encoding a protein that modulates the activity of the regulatory element may be identified by (i) inserting a nucleic acid that includes the second positive selection marker and a segment of the eukaryotic genome flanking the second integrated cassette into yeast cells under conditions that allow the selection of yeast cells expressing the second positive selection marker under the control of the yeast promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected yeast cells, and (iii) sequencing the amplified segment.

In a related aspect, the invention features another method for identifying a nucleic acid of interest that encodes a protein that modulates the activity of a regulatory element in a cell. This method involves inserting a first vector including a first cassette having a positive selection marker, a negative selection marker, and a recombinase signal sequence into eukaryotic cell under conditions that result in the integration of the first cassette into the genome of the cells. A second vector including a second cassette that includes a recombinase signal sequence, a nucleic acid segment encoding a transactivator polypeptide, and a reporter gene is inserted into the cells under conditions that result in

recombination between the recombinase signal sequence in the second vector and the recombinase signal sequence integrated into the genome of the cells. This recombination results in the integration of the second cassette into the genome of the cells such that the reporter gene is operably linked to a regulatory element in at least one cell. The

5 regulatory element may be an endogenous regulatory element, or the regulatory element may be a regulatory element of interest from the first or second vector. A third vector including a third cassette having a promoter operably linked to a responsive element that is responsive to the transactivator polypeptide is inserted into the cells. This step results in integration of the third cassette into the genome of the cells such that the promoter is

10 operably linked to an endogenous nucleic acid of interest encoding a protein that modulates (*i.e.*, increases or decreases) the activity of the regulatory element in at least one cell. The cells that have an altered level of reporter gene expression under conditions that activate the transactivator polypeptide are selected. Desirably, the nucleic acid of interest is identified from at least one selected cell. The nucleic acid of interest is

15 determined to encode a protein that activates the regulatory element if the cells have increased reporter gene expression under conditions that activate the transactivator polypeptide. Alternatively, the nucleic acid of interest is determined to encode a protein that inactivates the regulatory element if the cells have decreased reporter gene expression under conditions that activate the transactivator polypeptide. Desirably, the transactivator

20 polypeptide is tTA, and the responsive element comprises a tetracycline responsive element. Exemplary recombinase signal sequences include LoxP sites, Lox 511 sites, and any other recombinase signal sequence described herein. In various embodiments, the first and/or second vector include two recombinase signal sequences, such as two LoxP sites. In various embodiments, the first vector, second vector, third vector, or another

25 vector inserted into the cells encodes a recombinase that recognizes the recombinase signal sequence. In desirable embodiments, the recombinase signal sequence(s) in the first vector are identical to those in the second vector. In various embodiments, the first or second vector contains a regulatory element that was identified as being regulated by a stimulatory agent of interest. For example, the methods of the invention may be used to

30 identify an endogenous regulatory element that is regulated by a stimulatory agent, and then this regulatory element may be cloned into the first or second vector using standard

methods and used to identify endogenous nucleic acids encoding proteins that modulate the regulatory element. In other embodiments, the third vector further includes a positive selection marker which is integrated into the genome of the cells. In still other embodiments, the positive selection marker in the third vector (denoted the second

5 positive selection marker) is operably linked to a prokaryotic promoter in the third cassette that integrates into the genome of the eukaryotic cells. In this case, the nucleic acid encoding a protein that modulates the activity of the regulatory element may be identified by (i) inserting a nucleic acid that includes the second positive selection marker and a segment of the eukaryotic genome flanking the third integrated cassette into

10 bacterial cells under conditions that allow the selection of bacterial cells expressing the second positive selection marker under the control of the prokaryotic promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected bacterial cells, and (iii) sequencing the amplified segment. In yet other embodiments, the positive selection marker in the third vector is operably linked to a yeast promoter in the

15 third cassette that integrates into the genome of the eukaryotic cells. In this case, the nucleic acid encoding a protein that modulates the activity of the regulatory element may be identified by (i) inserting a nucleic acid that includes the second positive selection marker and a segment of the eukaryotic genome flanking the third integrated cassette into yeast cells under conditions that allow the selection of yeast cells expressing the second

20 positive selection marker under the control of the yeast promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected yeast cells, and (iii) sequencing the amplified segment.

In a related aspect, the invention features yet another method for identifying a nucleic acid of interest that encodes a protein that modulates the activity of a regulatory

25 element in a cell. This method involves inserting a first vector including a first cassette having a positive selection marker, a negative selection marker, a reporter gene, and a recombinase signal sequence into eukaryotic cell under conditions that result in the integration of the first cassette into the genome of the cells. A second vector including a second cassette that includes a recombinase signal sequence, and a nucleic acid segment

30 encoding a transactivator polypeptide, is inserted into the cells under conditions that result in recombination between the recombinase signal sequence in the second vector

and the recombinase signal sequence integrated into the genome of the cells. This recombination results in the integration of the second cassette into the genome of the cells such that the reporter gene is operably linked to a regulatory element in at least one cell. The regulatory element may be an endogenous regulatory element, or the regulatory element may be a regulatory element of interest from the first or second vector. A third vector including a third cassette having a promoter operably linked to a responsive element that is responsive to the transactivator polypeptide is inserted into the cells. This step results in integration of the third cassette into the genome of the cells such that the promoter is operably linked to an endogenous nucleic acid of interest encoding a protein that modulates (*i.e.*, increases or decreases) the activity of the regulatory element in at least one cell. The cells that have an altered level of reporter gene expression under conditions that activate the transactivator polypeptide are selected. Desirably, the nucleic acid of interest is identified from at least one selected cell. The nucleic acid of interest is determined to encode a protein that activates the regulatory element if the cells have increased reporter gene expression under conditions that activate the transactivator polypeptide. Alternatively, the nucleic acid of interest is determined to encode a protein that inactivates the regulatory element if the cells have decreased reporter gene expression under conditions that activate the transactivator polypeptide. Desirably, the transactivator polypeptide is tTA, and the responsive element comprises a tetracycline responsive element. Exemplary recombinase signal sequences include LoxP sites, Lox 511 sites, and any other recombinase signal sequence described herein. In various embodiments, the first and/or second vector include two recombinase signal sequences, such as two LoxP sites. In various embodiments, the first vector, second vector, third vector, or another vector inserted into the cells encodes a recombinase that recognizes the recombinase signal sequence. In desirable embodiments, the recombinase signal sequence(s) in the first vector are identical to those in the second vector. In various embodiments, the first or second vector contains a regulatory element that was identified as being regulated by a stimulatory agent of interest. For example, the methods of the invention may be used to identify an endogenous regulatory element that is regulated by a stimulatory agent, and then this regulatory element may be cloned into the first or second vector using standard methods and used to identify endogenous nucleic acids encoding proteins that modulate

the regulatory element. In other embodiments, both the first and second vectors have a reporter gene. In yet other embodiments, the third vector further includes a positive selection marker which is integrated into the genome of the cells. In still other

5 positive selection marker) is operably linked to a prokaryotic promoter in the third cassette that integrates into the genome of the eukaryotic cells. In this case, the nucleic acid encoding a protein that modulates the activity of the regulatory element may be identified by (i) inserting a nucleic acid that includes the second positive selection marker and a segment of the eukaryotic genome flanking the third integrated cassette into

10 bacterial cells under conditions that allow the selection of bacterial cells expressing the second positive selection marker under the control of the prokaryotic promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected bacterial cells, and (iii) sequencing the amplified segment. In yet other embodiments, the positive selection marker in the third vector is operably linked to a yeast promoter in the

15 third cassette that integrates into the genome of the eukaryotic cells. In this case, the nucleic acid encoding a protein that modulates the activity of the regulatory element may be identified by (i) inserting a nucleic acid that includes the second positive selection marker and a segment of the eukaryotic genome flanking the third integrated cassette into yeast cells under conditions that allow the selection of yeast cells expressing the second

20 positive selection marker under the control of the yeast promoter, (ii) amplifying the segment of the eukaryotic genome that is inserted into the selected yeast cells, and (iii) sequencing the amplified segment.

In yet another aspect, the invention provides a method for treating, preventing, or stabilizing a disease that is mediated by, or associated with, a stimulatory agent. This

25 method involves identifying a cell containing a regulatory element that is regulated by a stimulatory agent, selecting a compound that modulates the regulatory element or that modulates a protein which regulates the regulatory element, and administering the compound to a mammal having a disease or condition associated with the stimulatory agent or having an increased risk for the disease or condition. The cells that are regulated

30 by stimulatory agent of interest may be identified using any of the methods of the invention.

In particular embodiments of the above aspect, the method involves inserting a vector which has a cassette including a positive selection marker, a negative selection marker, and a reporter gene into eukaryotic cells under conditions that result in the integration of the cassette into the genome of the cells such that the reporter gene is operably linked to a regulatory element in at least one cell. Cells in which expression of the reporter gene is specifically modulated by the stimulatory agent are selected. A compound that increases or decreases the effect of the stimulatory agent on the expression of the reporter gene is selected and administered to a mammal having a disease associated with the stimulatory agent. If the stimulatory agent is associated with an increased risk for the disease or is associated with increased severity of the disease, the administered compound preferably inhibits the ability of the stimulatory agent to modulate the expression of the reporter gene. Conversely, if the stimulatory agent is associated with a decreased risk for the disease or is associated with decreased severity of the disease, the administered compound preferably enhances the ability of the stimulatory agent to modulate the expression of the reporter gene.

In another embodiment of the above aspect, the method involves inserting a vector which has a cassette including a positive selection marker, a negative selection marker, and a nucleic acid segment encoding a transactivator polypeptide into eukaryotic cells under conditions that result in the integration of the cassette into the genome of the cells such that the nucleic acid segment encoding a transactivator polypeptide is operably linked to a regulatory element in at least one cell. Cells in which expression of the transactivator polypeptide is specifically modulated by the stimulatory agent are selected. A compound that increases or decreases the effect of the stimulatory agent on the expression of the transactivator polypeptide is selected and administered to a mammal having a disease associated with the stimulatory agent. If the stimulatory agent is associated with an increased risk for the disease or is associated with increased severity of the disease, the administered compound preferably inhibits the ability of the stimulatory agent to modulate the expression of the transactivator polypeptide. Conversely, if the stimulatory agent is associated with a decreased risk for the disease or is associated with decreased severity of the disease, the administered compound preferably enhances the

ability of the stimulatory agent to modulate the expression of the transactivator polypeptide.

In still another aspect, the invention features a nucleic acid including a positive selection marker, a negative selection marker, and a reporter gene. In particular  
 5       embodiments, the nucleic acid includes, in 5' to 3' order, a splice acceptor, a cassette including in any order a negative selection marker and a positive selection marker, a translation stop sequence, an internal ribosome entry site, a reporter gene, a translation stop sequence, and a polyadenylation signal. In another embodiment, the nucleic acid includes, in 5' to 3' order, a splice acceptor, a cassette including in any order a negative  
 10       selection marker and a reporter gene, a translation stop sequence, a promoter, a positive selection marker, a translation stop sequence, and a polyadenylation signal. In other embodiments, the reporter gene is not operably linked to a promoter in the nucleic acid. In this embodiment, the nucleic acid may be inserted in a cell such that the reporter gene is operably linked to an endogenous promoter. In other embodiments, the nucleic acid  
 15       also includes a nucleic acid segment encoding a transactivator polypeptide or also includes one or more recombinase signal sequences (*e.g.*, LoxP sites).

In yet another aspect, the invention features a nucleic acid including a splice acceptor site and including a bacterial promoter operably linked to a positive selection marker or a reporter gene. In various embodiments, the nucleic acid also includes a  
 20       negative selection marker which may or may not be operably linked to the bacterial promoter. In other embodiments, the nucleic acid also includes a translation stop sequence, an internal ribosome entry site, a reporter gene, a translation stop sequence, and a polyadenylation signal. In particular embodiments, the nucleic acid includes, in 5' to 3' order, a splice acceptor, a cassette including in any order a negative selection marker and  
 25       a positive selection marker such that the positive selection marker is operably linked to a bacterial promoter, a translation stop sequence, an internal ribosome entry site, a reporter gene, a translation stop sequence, and a polyadenylation signal. In another embodiment, the nucleic acid includes, in 5' to 3' order, a splice acceptor, a cassette including in any order a negative selection marker and a reporter gene, a translation stop sequence, a  
 30       bacterial promoter operably linked to a positive selection marker, a translation stop sequence, and a polyadenylation signal. In other embodiments, the positive selection

marker is operably linked to the bacterial promoter, and the reporter gene is not operably linked to a promoter in the nucleic acid. In this embodiment, the nucleic acid may be inserted in a cell such that the reporter gene is operably linked to an endogenous promoter. In other embodiments, the nucleic acid also includes a nucleic acid segment encoding a transactivator polypeptide or also includes one or more recombinase signal sequences (*e.g.*, LoxP sites). In still other embodiments, the nucleic acid includes a region of a eukaryotic genome, such as a region containing all or part of a gene or a regulatory element of interest or a region flanking a gene or a regulatory element of interest. Such nucleic acids enable bacterial cells to be used to facilitate the identification of trapped eukaryotic regulatory elements or genes of interest, as described herein.

In a related aspect, the invention features a nucleic acid including a positive selection marker, a negative selection marker, and a nucleic acid segment encoding a transactivator polypeptide. In various embodiment, the nucleic acid also includes one or more recombinase signal sequences (*e.g.*, LoxP sites). In other embodiments, the nucleic acid segment encoding the transactivator polypeptide is not operably linked to a promoter in the nucleic acid.

In another related aspect, the invention provides a nucleic acid including a positive selection marker, a negative selection marker, and one or more recombinase signal sequences (*e.g.*, LoxP sites).

In still another aspect, the invention features a nucleic acid including, in 5' to 3' sequence, an internal ribosome entry site, a nucleic acid segment encoding a transactivator polypeptide, a translation stop sequence; an internal ribosome entry site, a reporter gene; a translation stop sequence, and a polyadenylation signal. The nucleic acid may also include a recombinase signal sequence (*e.g.*, a LoxP site). In another aspect, the invention provides a nucleic acid having a functional splice acceptor, a translation stop sequence, an internal ribosome entry site, a promoterless negative selection marker, a translational stop sequence, a polyadenylation signal, a promoter, positive selection marker, a translational stop sequence, and a polyadenylation signal. In other embodiments, the nucleic acid also includes an internal ribosome entry site, a nucleic acid segment encoding a transactivator polypeptide, a translation stop sequence, an internal ribosome entry site, a reporter gene, and a translation stop sequence. In yet other

embodiments, the nucleic acid also includes an internal ribosome entry site, a nucleic acid segment encoding a transactivator polypeptide, and a translation stop sequence.

In another aspect, the invention features a vector, such as a retroviral vector that contains one or more nucleic acids of the invention. The vector may optionally include  
 5 an integration sequence. In particular embodiments, the retroviral vector is a replication deficient viral vector, such as a SIN virus viral vector that contains a mutation in the 3' LTR.

In yet another aspect, the invention features a cell (*e.g.*, a eukaryotic or prokaryotic cell) containing a vector or nucleic acid of the invention. The cell may be  
 10 responsive to only one or to more than one stimulatory agent. Exemplary cells contain (i) a first nucleic acid which includes a positive selection marker, a negative selection marker, and a nucleic acid segment encoding a transactivator polypeptide and (ii) a second nucleic acid which includes a promoter operably linked to a responsive element that is responsive to the transactivator polypeptide. In particular embodiments, the first  
 15 nucleic acid also includes a reporter gene, or the second nucleic acid also includes a positive selection marker.

In a related aspect, the invention features a library of two or more cells (*e.g.*, eukaryotic or prokaryotic cells) containing a vector or nucleic acid of the invention. In particular embodiments, the library of cells contains at least 5, 10, 20, 50, 100, 500, 1000,  
 20 50000, or more cells containing different trapped regulatory elements (*e.g.*, different endogenous regulatory elements operably linked to a positive selection marker or reporter gene in a construct integrated into the genome of the cells) or different trapped genes (*e.g.*, different endogenous genes operably linked downstream of a promoter in a construct integrated into the genome of the cells). Desirably, the library includes cells  
 25 that are responsive to one or more stimulatory agents of interest. In other embodiments, the library includes 5, 10, 20, 50, 100, 500, 1000, 50000, or more different cells that are each responsive to a different stimulatory agent of interest.

In still another aspect, the invention features a screening method for selecting compounds that modulate the activity of a stimulatory agent of interest. This method  
 30 includes contacting one or more cells of the invention that have a specific response to the stimulatory agent with one or more candidate compounds and the stimulatory agent. The

candidate compounds which modulate (*i.e.*, increase or decrease) the response to the stimulatory agent are selected.

In yet another aspect, the invention features a method for determining whether a candidate compound modulates the activity of a regulatory element of interest. The method includes contacting one or more cells of the invention that have the regulatory element of interest operably linked to a positive selection marker, reporter gene, or nucleic acid segment encoding a transactivator polypeptide with one or more candidate compounds. A candidate compound which modulates the expression of the positive selection marker, reporter gene, or nucleic acid segment encoding a transactivator polypeptide is selected, thereby selecting a candidate compound which modulates the activity of the regulatory element of interest. In particular embodiments, the modulation (*i.e.*, increase or decrease) of the activity of the regulatory element of interest is associated with adverse side-effects of the candidate compound *in vivo*. In this case, the candidate compound is desirably eliminated from drug development due to the potential adverse side-effects (*e.g.*, drug toxicity) of the candidate compound when administered to a mammal (*e.g.*, a human). For example, a candidate compound that activates a regulatory element operably linked to a gene encoding an mRNA or a protein involved in a pathway associated with adverse side-effects is desirably eliminated from further drug development. In other embodiments, the method is used to determine whether particular combinations of two or more candidate compounds are likely to be associated with adverse side-effects when administered together (*e.g.*, sequentially or concurrently) to a mammal. For example, a combination of candidate compounds that activates a regulatory element operably linked to a gene encoding an mRNA or a protein involved in a pathway associated with adverse side-effects is desirably eliminated from further drug development. In yet other embodiments, the method is performed prior to animal model studies or human clinical trials of the candidate compound or the combination of candidate compounds to determine whether or not the candidate compound(s) are likely associated with adverse side-effects prior to further drug development.

In various embodiments of any of the aspects of the invention, the cells are mast cells, stem cells, epithelial cells, fibroblast cells, cancer cells, lymphocytes, and liver cells. Other exemplary cells include cells from tumor cell lines from cancers of one of

the following cell types: rectum, colon, ovary, prostate, pancreas, mammary gland, lung, ovary, kidney, cervix, tongue, thyroid, T lymphocyte, B lymphocyte, adenocarcinoma, small cell lung, burkitt's lymphoma, adenosquamous carcinoma, adrenocortical carcinoma, alveolar cell carcinoma, or hodgkin's lymphoma. An example of a eukaryotic genome is the genome of a mammalian cell. Suitable stimulatory agents include cytokines, growth factors, ligands, polypeptides, growth factors, antibodies, and chemical agents. Exemplary stimulatory agents include stem cell factor, IL-1, IL-3, IL-2, IL-6, IL-8, IL-18, IgE, Fibroblast Growth Factors (FGFs), FGF-1, FGF-2, FGF-3, transforming growth factor  $\alpha$  (TGF- $\alpha$ ), TGF- $\beta$ , TNF- $\beta$ , TNF- $\alpha$ , VEGF, leptin, epidermal growth factor (EGF), platelet-derived growth factor (PDGF), insulin, insulin-like growth factor-I& II, interferon- $\gamma$  (IFN- $\gamma$ ), estrogen, testosterone, and colony stimulating factors (CSFs). It is also contemplated that the stimulatory agent controls the expression of an exogenous gene that is inserted into the cells. For example, the stimulatory agent (*e.g.*, a chemical agent) can activate the promoter operably linked to an exogenous gene that encodes a protein which modulates the activity of an endogenous regulatory element of interest in the cells. In various embodiments, a protein activates the activity of an endogenous regulatory element of interest by inducing an activator or by inhibiting a repressor of the regulatory element. In other embodiments, the protein inhibits the activity of an endogenous regulatory element of interest by inhibiting an activator or by activating a repressor of the regulatory element. In other embodiments, the nucleic acid, cassette, vector, or cell includes a prokaryotic promoter (*e.g.*, a bacterial promoter) or yeast promoter operably linked to a positive selection marker or reporter gene.

Desirable reporter genes encode an enzyme, such as secreted alkaline phosphatase,  $\beta$ -galactosidase, luciferase, and green fluorescent protein. In any of the above aspects, a nucleic acid segment encoding a single protein that has both positive selection traits and negative selection traits may be used as the positive and negative selection markers. In other embodiments, the negative selection marker and the positive selection marker encode different proteins. In still other embodiments, the reporter gene is different from the positive selection marker and/or the negative selection marker. Exemplary negative selection markers include nucleic acid segments encoding Hp<sub>r</sub>rt, gpt, HSV-tk, diphtheria toxin, ricin toxin, or cytosine deaminase. Exemplary positive

selection markers include nucleic acid segments encoding proteins conferring neomycin resistance, hygromycin resistance, histidinol resistance, xanthine utilization, Zeocin resistance, or bleomycin resistance. Examples of internal ribosome entry sites include mammalian, picornavirus, and polio internal ribosome entry sites.

5

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a schematic diagram of a nucleic acid construct of this invention for use in preparing an induction gene trap vector. The following components are illustrated in the diagram: an internal ribosome entry site (IRES), a promoterless protein coding sequence coding a tetracycline regulator protein ("TetOn/Off"), an internal ribosome entry site, and a secreted alkaline phosphatase ("SEAP").

10

Fig. 2 is a schematic diagram of a vector including the nucleic acid construct of Fig. 1 which also contains, upstream of the construct, the following additional components: a functional splice acceptor (SA), a translation stop sequence ("STOP"), an IRES site, and a promoterless protein coding sequence coding TK-ZEO; and a polyadenylation signal ("pA") downstream of the nucleic acid sequence of Fig. 1.

15

Fig. 3 is a schematic diagram of another alternate vector for use in the invention which includes the nucleic acid construct of Fig. 1 and, upstream of the construct, a functional splice acceptor (SA), and a translation stop sequence (STOP), and downstream of the construct, an IRES site, a TK coding sequence, a phosphoglycerate kinase promoter ("PKG"), a ZEO coding sequence under the transcriptional control of the PKG promoter, and a polyadenylation signal (pA).

20

Fig. 4 is a schematic diagram of a nucleic acid construct of this invention for use in a vector in conjunction with the induction trap vectors of FIGS. 2 and 3. The following components are illustrated in the diagram: a minimal promoter sequence containing a tetracycline responsive element ("TRE<sub>pcmv</sub>"), a promoterless protein coding sequence encoding NEO, an IRES sequence, and a splice donor ("SD").

25

Fig. 5 is a schematic diagram illustrating the use of a vector containing the construct of Fig. 1 and a vector containing the nucleic acid construct of Fig. 4 which can be integrated into the genomic loci to select genes that directly or indirectly regulate the

30

genomic loci. A putative gene transcribed by the vector containing the nucleic acid construct of Fig. 4 is also shown.

Fig. 6 is a schematic representation illustrating the preparation of a ligand dependent cell from a selected cell line by contacting the cell with the transfection vector of Fig. 2, a physiological stimuli, and positive and negative selection drugs. The selection of specific cells which are activated or turned on by the physiological stimuli, and cells which are inactivated or turned off by the physiological stimuli, are illustrated in the left and right branches of the diagram, respectively.

Fig. 7 is a schematic representation illustrating an alternate method for the preparation of a ligand dependent cell from a selected cell line by contacting the cell with the transfection vector of Fig. 3, a physiological stimuli, and positive and negative selection drugs. The selection of specific cells which are activated or turned on by the physiological stimuli, and cells which are inactivated or turned off by the physiological stimuli, is illustrated in the left and right branches of the diagram, respectively. The production of SEAP is measured as an indicator of the response of the vector to the stimuli.

Fig. 8A is a schematic illustration of an induction trap vector. This vector includes a functional splice acceptor (SA), a translation stop sequence (STOP), an IRES site, a TK coding sequence, a ZEO coding sequence, a STOP sequence, a LoxP site, a IRES site, a SEAP coding sequence, a STOP sequence, a polyadenylation signal (PolyA), and a LoxP site. Other vectors that may be used in the methods of the invention include the corresponding induction trap vectors that contain only one LoxP site or that lack LoxP sites.

Fig. 8B is a schematic illustration of an exchange cassette that is used to replace the region of the induction trap vector of Fig. 8A that is flanked by LoxP sites, as described in Example 10. This cassette includes a LoxP site, an IRES site, a promoterless sequence encoding a tetracycline regulator protein ("teton/off"), a translation stop sequence (STOP), an IRES site, a  $\beta$ -galactosidase coding sequence (b-gal), a STOP sequence, a polyadenylation signal (PolyA), and a LoxP site. Other exchange cassettes that may be used in the methods of the invention include the corresponding cassettes with only one LoxP site. The LoxP site may be located in any part of the cassette.

Fig. 9A is a schematic illustration of a vector of the present invention. This vector contains a prokaryotic promoter operably linked to a positive selection marker (*e.g.*, zeocin). This exemplary vector contains a functional splice acceptor (EN-2 SA), a translation stop sequence (STOP), an IRES site, a prokaryotic promoter, a negative selection marker (*e.g.*, TK coding sequence), a positive selection marker (*e.g.*, a ZEO coding sequence), another STOP sequence, a LoxP site, an IRES site, a reporter gene (*e.g.*, a SEAP coding sequence), a STOP sequence, a polyadenylation signal (PolyA), and a LoxP site. The ClaI site represents a possible location of a unique restriction site in the vector. One skilled in the art would readily appreciate that the components of this vector may be present in different locations or different 5' to 3' arrangements. For example, the prokaryotic promoter may alternatively be located upstream of the first IRES site or between the TK coding sequence and the Zeo coding sequence. As noted above, the Zeo coding sequence may also be located upstream, instead of downstream, of the TK coding sequence. In some methods of the invention, the vector may lack the first LoxP site, the second IRES site, the SEAP coding sequence, the third STOP sequence, and/or the second LoxP site. Other vectors of the present invention contain a yeast promoter instead of the prokaryotic promoter in any of the vectors described above.

Fig. 9B is the polynucleotide sequence of an exemplary prokaryotic promoter, the T7 promoter (SEQ ID NO: 5). "RBS" denotes a ribosome binding site. Any other prokaryotic promoter or any yeast promoter may also be used in the nucleic acids, vectors, cells, and methods of the invention.

Fig. 10 is a schematic illustration of the uses of the cells of the invention to identify ligand specific pathways, redundant pathways, and pathways associated with toxic effects *in vivo*. This information is useful in the characterization of candidate drug products and the prediction of adverse side-effects caused by these products.

Fig. 11 is a schematic illustration of the use of the methods described herein to isolate EL4 or NIH3T3 fibroblast cells activated by  $\text{TNF}\alpha$  or  $\text{IL-1}\beta$ .

Fig. 12A is a table confirming that the reporter gene (SEAP) that integrated into the genome of cells was integrated under the control of a regulatory element responsive to  $\text{TNF}\alpha$  or  $\text{IL-1}\beta$ . Fig. 12B is a picture of a southern blot generated using a probe to the

TK/Zeo selection markers in the integrated construct to confirm the integration of the construct in some of the selected NIH3T3 cell lines.

Fig. 13A is a schematic illustration of the identification of cells responsive to a single or multiple ligands. Fig. 13B is a bar graph illustrating the level of responsiveness of selected cell clones to IL-1 $\beta$ , TNF $\alpha$ , and IL-6.

Figs. 14A and 14B are a set of bar graphs illustrating the level of responsiveness of selected clones to IL-1 $\beta$ , TNF $\alpha$ , PMA, and IL-10. Fig. 14C is a bar graph illustrating the level of responsiveness of selected clones to IL-1 $\beta$ , TNF $\alpha$ , SDF-1, MCP-1, and IL-10.

Fig. 15 is a graph illustrating the ability of the specific Cox-2 inhibitor, celecoxib, to inhibit the effect of IL-1 $\beta$  on SEAP reporter gene activity in selected NIH3T3 cells in a concentration dependent manner.

Fig. 16 is a graph illustrating the inability of celecoxib to significantly inhibit TNF $\alpha$ -induced SEAP reporter gene activity in selected EL-4 cells.

Figs. 17A and 17B are a set of bar graphs illustrating the level of responsiveness to various ligands and ligand combinations in clone C-5 and clone PD6.

Fig. 18A is a graph illustrating the ability of the MEK inhibitor U0126 to inhibit the effect of IL-1 $\beta$  on SEAP reporter gene activity in selected NIH3T3 cells in a concentration dependent manner. As illustrated in Fig. 18B, cyclosporin A had a much smaller effect on SEAP activity in this assay.

## DETAILED DESCRIPTION OF THE INVENTION

The methods of this invention utilize cells, vectors, and stimulatory agents to generate cell lines, and to identify gene targets and regulatory elements which are useful for the selection of therapeutic agents from a library of drug candidates.

The cells which are useful in this invention are eukaryotic cells, preferably mammalian cells, and more preferably human cells. The eukaryotic cells are capable of differentiating into specific cell or tissue types, including both plant and animal cells and tissues. Particularly suitable are totipotent cells, such as stem cells, as well as mast cells, endothelial cells, epithelial cells, cancer cells, lymphocytes, and liver cells.

A reporter element useful in the nucleic acid constructs and vectors of this invention are elements which express indicators in cells which are capable of being

detected using physical, chemical, or optical means. The detection can be visual, instrument assisted or completely automated. Suitable reporter elements include enzymes, such as secreted alkaline phosphatase, luciferase, and green fluorescent protein. Enzymes which emit fluorescence can be detected using a luminometer.

5           An “induction gene trap vector” means a vector containing elements that allow for selection and insertion of the trap vector in an operably linked manner into an intron sequence of a regulated genomic loci of a cell, by techniques well known in the art, such as transfection, transduction, and the like, resulting in the transformation and integration of the vector into the genome. The induction gene trap vector contains a marker gene  
10       sequence expressing selection traits, such as positive and negative selection traits.

          The induction gene trap vector may be “promoterless,” which means that the marker genes are not under the control of a promoter within the vector (although the vector may contain promoters which do not regulate these elements). In the case of a  
15       regulatory elements or factors in the genome which respond to one or more exogenous stimulatory agents externally introduced into the cell.

          Alternatively, the vector may contain a promoter for at least one component of the marker gene sequence, such as the PGK (phosphoglycerate kinase) promoter for the neo positive selection marker, as described in Mainguy *et al.*, *Nature Biotechnology*, Vol. 18,  
20       pages 746-749 (2000). A vector containing a promoter for one such component, but lacking promoters for sequences expressing other selectable traits and for reporter sequences, is also included within the scope of this invention.

          Elements or sequences in a vector which are “operably linked,” and vectors which are “operably integrated” into a genome, refer to nucleotide sequences which are linked,  
25       whether to encode an mRNA transcript of a desired gene product, or for regulatory control. “Operably linked” can also mean that selectable marker, transactivator, and reporter genes are encoded by the same transcription unit.

          A “splice acceptor (“SA”), or functional splice acceptor, refers to a consensus sequence that permits the construct or vector to be processed such that it is included in a  
30       mature, biologically active mRNA, provided that it is integrated in an active chromosomal locus and transcribed as a contiguous part of the premessenger RNA of the

chromosomal locus. Splice acceptors typically include the 3' end of an intron and the 5' end of an exon, while a splice donor ("SD") typically includes the 5' end of an exon and the 3' end of an intron. Examples of these elements, as well as other gene elements used to prepare gene trap vectors, can be found in published patent application

- 5 PCT/CA98/00667, Alberts *et al.*, *Molecular Biology of the Cell*, page 373 (1994), and U.S. Patent no. 5,922,601, the disclosures of which are incorporated herein by reference thereto in their entirety.

A translation stop sequence, or "STOP," is a sequence that codes for translation stop codons in three different reading frames. The STOP sequence causes truncation of  
10 peptide chains encoded by exons upstream of the vector at the chromosomal locus and prevents the translational reading frame from proceeding into the selectable marker gene, thereby preventing translating in a non-sense reading frame.

An internal ribosome entry site, or "IRES," as used herein, is an element which permits attachment of a downstream coding region or open reading frame with a  
15 cytoplasmic polysomal ribosome to initiate translation thereof in the absence of internal promoters. An IRES is included in the construct to initiate translation of selectable marker protein coding sequences. The encephalomyocarditis virus IRES is one such IRES which is suitable for use in this invention.

A "marker" refers to nucleotide sequences in vectors or genes encoding  
20 polypeptides or proteins which can be used to distinguish cells expressing the protein from those not expressing the protein. Marker genes can be detected using a variety of means and include selectable markers and assay markers. Selectable markers are genetic elements which can be selected or screened for when integrated into the genome or genomic loci of a cell. Selectable markers include markers having selection traits, such as  
25 drug resistant markers, antigenic markers, adherence markers, and the like. Examples of antigenic markers include those useful in fluorescence-activated cell sorting. Examples of adherence markers include receptors for adherence ligands that allow selective adherence. Other selection markers include a variety of gene products that can be detected in experimental assay protocols, such as marker enzymes, amino acid sequence  
30 markers, cellular phenotypic markers, nucleic acid sequence markers, and the like.

The selectable markers also include markers with both negative and positive selection traits. In general, positive selection refers to the isolation of cells that express the marker gene, and negative selection refers to the isolation of cells that do not express the marker gene. In various embodiments, the expression of a negative selection marker leads to the selective elimination or death of cells containing the marker. A single gene or multiple genes can be used for positive and negative selection. Gene sequences which express a fusion protein having both positive and negative selection traits are preferred. As a specific example, a fusion protein can be expressed by a gene sequence encoding the negative selection marker Tk (thymidine kinase) and the positive selection marker neo (neomycin phosphotransferase). Details concerning gene markers having positive and/or negative selection traits and additional examples of selectable markers can be found in U.S. Patent No. 5,922,601; filed September 16, 1996; issued July 13, 1999, the disclosure of which is incorporated by reference thereto in its entirety. Any of these selectable markers may also be used in the nucleic acid constructs and methods of the present invention.

A “transactivator” and a “transactivator polypeptide” are nucleic acid sequences and polypeptides, respectively, that transcribes, or causes the transcription of, a protein which effects the regulation of a genomic loci. Examples of transactivator polypeptides include transcription factors and growth factors. Other exemplary transactivator polypeptides include molecules involved in a signaling pathway. The transactivator polypeptides may directly or indirectly activate the transcription of a gene. For example, a transactivator polypeptide may directly bind a regulatory element; such as an enhancer, transcription factor binding site, or promoter; and activate the transcription of a gene downstream of the regulatory element. Alternatively, a transactivator polypeptide may activate another polypeptide that directly or indirectly activates transcription of the gene.

A “regulator unit or regulator protein” is a transactivator polypeptide that binds regulatory elements which effect the regulation of a genomic loci. For example, the transactivator polypeptide may bind a regulatory element (such as a tetracycline responsive promoter) and activate the transcription of an endogenous gene that is downstream of the regulatory element. The protein encoded by this endogenous gene may then activate a

regulatory element of interest (such as an endogenous promoter or other regulatory element identified using an induction trap vector of the present invention).

A tetracycline regulator unit is an example of a transactivator regulatory sequence which expresses a protein ("tTA") activated or repressed by tetracycline. The tetracycline  
 5 regulator unit can be incorporated in a vector which acts in concert with a minimal promoter sequence containing tetracycline responsive elements, or "TRE<sub>Pcmv</sub>," which is present in a complementary vector. See U.S. Patent No. 5,464,758 and U.S. Patent No. 5,814,618, the disclosures of which are incorporated herein by reference in their entirety. This pair of vectors can be operably integrated into the genome of a cell. When the cell is  
 10 contacted with a stimulatory agent, the tetracycline regulator is turned on, causing the unit to generate a protein which binds to the minimal promoter sequence containing the tetracycline responsive element. This causes the minimal promoter to activate and induce transcription of genes downstream of the promoter. These transcribed genes can up regulate or down regulate the genomic loci, causing the tetracycline regulator unit to  
 15 express more protein, thereby activating the promoter to transcribe additional copies of the gene, and so on. Eventually, as a result of this feedback process, enough genetic material is generated to be detected, isolated and sequenced.

"5' RACE" cloning, as that expression is used herein, refers to 5' rapid PCR amplification of cDNA ends (RACE). This procedure is described in detail by Skarnes *et al.*, *Genes and Development*, 6, pages 903-918 (1992), the disclosure of which is  
 20 incorporated herein by reference in its entirety.

By "cassette" is meant a segment of a nucleic acid.

By "polypeptide" is meant a sequence of two or more covalently bonded naturally-occurring or modified amino acids. The terms "peptide," "polypeptide," and  
 25 "protein" are used interchangeably herein.

The method and vectors of this invention can be used to select cell lines and identify regulatory components which respond to stimulation from a selected stimulatory agent. This is accomplished by utilizing induction gene trap vectors to introduce specific polynucleotide sequences into genomic loci which respond to selected stimuli. While  
 30 certain specific nucleotides and vectors have been illustrated herein, this is done for convenience in understanding the invention only and is not intended to limit the scope of

the invention. Other vectors can be readily designed by those skilled in the art and advantageously used to practice the methods described herein.

In general, the induction trap vectors of this invention contain a transactivator gene or polypeptide coding sequence, and/or contain a reporter element or sequence. The reporter element is preferably a sequence encoding an enzyme which is capable of being detected. Suitable enzymes are well known in the art and include secreted alkaline phosphatase (SEAP), Luciferase<sup>TM</sup> and green fluorescent protein. Enzymes emitting light can be detected using, for instance, a fluorescent activated cell sorter or similar device.

One specific nucleic acid construct is shown in Fig. 1. This construct can be incorporated in an induction gene trap vector and used to transfect cells of interest. As shown in Fig. 1, some constructs which are operable in this invention include a cassette containing an internal ribosome entry site, a transactivator gene such as a promoterless protein coding sequence coding a tetracycline regulator protein, an internal ribosome entry site, and a reporter sequence such as secreted alkaline phosphatase. Obviously, other transactivator genes and reporter elements can be used in the construct in place of the specific components shown in the Fig. 1. Moreover, a translation stop sequence can be inserted between the tetracycline regulator unit and the IRES sequence, and a STOP sequence can be used at the end of the construct as well. The construct can be incorporated into a vector, such as a viral vector, for use in transfecting cells.

Another vector which is useful in this invention is illustrated in Fig. 2. The vector of Fig. 2 contains the following operably linked components: a functional splice acceptor; a translation stop sequence; an IRES site; and a promoterless protein coding sequence encoding a fusion protein having positive and negative selection traits, such as the gene encoding the fusion protein for the negative/positive selection polypeptide Tk-Zeo; an internal ribosome entry site; a gene marker such as a promoterless protein coding sequence encoding a tetracycline regulator protein, an internal ribosome entry site; a reporter sequence such as a sequence encoding secreted alkaline phosphatase; and a polyadenylation signal. Some components of this vector may be redundant depending on the particular uses of the vector. For instance, if the vector is used to select for a cell line responsive to a stimulatory agent, it may be possible to eliminate the reporter element,

and its associated IRES, depending on the particular selection protocol used in the gene trap procedure, as illustrated in FIGs. 6 and 7.

The vector illustrated in Fig. 2, Fig. 8A, or Fig. 9A can be used to generate cell lines using the procedure outlined in Fig. 6. As shown, cells are transformed, using  
 5 suitable techniques such as transfection or transduction, with the retroviral vectors of Fig. 2. A stimulatory agent, such as IL-3, and a selection drug, such as zeocin, are added to the cell culture medium. The live cells remaining in the culture medium are cells with the vector integrated into the genomic loci that are (1) turned on (activated) by the stimulatory agent and (2) contain activated housekeeping genes. The live cells are  
 10 separated from the medium and placed in a fresh medium with gancyclovir to eliminate the cells with active housekeeping genes. Cells that are turned on by the stimulating agent are identified and isolated.

Alternatively, the cells in Fig. 6 are transformed with the retroviral vectors of Fig. 2, Fig. 8A, or Fig. 9A. A selection drug (zeocin) is added to the medium, and those cells  
 15 remaining are cells containing housekeeping genes, and cells which are turned on in the absence of the stimulatory agent. A stimulatory agent, such as IL-3, and gancyclovir are added to fresh medium containing the activated cells, and cells which contain activated housekeeping genes are eliminated. The cells remaining are those cells which are turned off by the stimulatory agent.

20 Fig. 2 illustrates another vector which can be used to generated cell lines. This vector includes the following operably linked components in downstream sequence: a functional splice acceptor sequence, a translation stop sequence, an internal ribosome entry site, a transactivator gene such as a promoterless protein coding sequence coding a tetracycline regulator protein, an internal ribosome entry site, a reporter sequence such as  
 25 secreted alkaline phosphatase, an IRES site, a coding sequence encoding TK, a phosphoglycerate kinase promoter ("PKG"), a Zeocin coding sequence under the transcriptional control of the PKG promoter, and a polyadenylation signal (pA).

The vector illustrated in Fig. 3, Fig. 8A, or Fig. 9A can be used to generate cell lines using the procedure outlined in Fig. 7. As shown in Fig. 7, cells are transformed  
 30 with the induction gene trap vector shown in Fig. 3, Fig. 8A, or Fig. 9A and a selection drug (*e.g.*, zeocin) is added to the medium to eliminate cells which do not have the vector

integrated into the genomic loci. The medium is changed, and gancyclovir is added to eliminate cells containing active housekeeping genes. A stimulatory agent is added, and those cells which respond to the agent are selected based on the amount of secreted alkaline phosphatase produced by the cells. These are cells which are activated by the stimulatory agent.

Alternatively, the cells in Fig. 7 are transformed with the retroviral vector of Fig. 3, and a selection drug (zeocin) is added to the medium. To eliminate cells which do not have the vector integrated into the genome. The medium is changed, and gancyclovir and a stimulatory agent are added to eliminate cells which contain housekeeping genes. Cells which are turned off by the stimulatory agent are selected by measuring the amount of secreted alkaline phosphatase produced by the cells in the absence of the stimulatory agent.

Fig. 4 illustrates a vector which can be used in combination with a vector containing the nucleic acid sequence shown in the shaded area depicted in Fig. 2 or Fig. 3 to identify a gene which is capable of up regulating or down regulating the genomic loci of a cell which responds to a stimulatory agent, such as the cells identified in FIGs. 6 or 7. The vector of Fig. 4 includes the following components operably linked in downstream sequence: a minimal promoter sequence containing a tetracycline responsive element; a promoterless protein coding sequence encoding Neo; an IRES sequence; and a splice donor. The shaded area in FIGs. 2 or 3 contains the following components in downstream sequence: an IRES sequence, a Tet On/Off sequence, an IRES sequence, and an SEAP expression sequence. Alternatively, a region of the induction trap vector of Fig. 8A may be replaced with the exchange cassette of Fig. 8B to generate a vector 5 that includes an IRES sequence, a TetOn/Off sequence, a STOP sequence, an IRES sequence, a  $\beta$ -galactosidase expression sequence, a STOP sequence, and a polyadenylation sequence. The secondary cell infection procedure is illustrated in Fig. 5.

As shown in Fig. 5, cell 1 having a nucleus 2 is transfected with vectors 4 and 5, and the vectors are integrated into the genomic loci 3. Vector 4, which can be the vector of Fig. 4, and vector 5, which can be a vector containing the nucleic acid sequence shown as the shaded area in FIGs. 2 or 3. are transfected and integrated into the genomic loci 3 of cell 1. Cell 1 can be a cell of the type depicted in Fig. 6 or Fig. 7. The integrated

vectors act in a complementary fashion to cause an increase in the expression of a gene downstream of the integration site of the transfected vector 4, and the expression of the tetracycline regulator protein coded by vector 5. This occurs as a result of the activation of the tetracycline regulator unit that transcribes protein 6 (tTA), which is a protein

5 activated or repressed by tetracycline. The tTA protein 6 binds to the minimal promoter sequence containing the tetracycline responsive element in vector 4, activating the TRE<sub>Pcmv</sub> promoter and transcribing additional protein 7 (protein X) of the downstream gene. The protein 7 transcribed by the downstream gene up regulates or down regulates the genomic loci 3, causing increased expression of the tetracycline regulator unit in

10 vector 5, thereby activating the TRE<sub>Pcmv</sub> promoter in vector 4 to transcribe additional protein 7 from the downstream gene. This process repeats itself in a continuous loop until sufficient protein 7 is transcribed to permit the collection and identification of protein 7 or its mRNA. This allows the corresponding gene to be identified and characterized. As shown in Fig. 5, the gene transcription process can be monitored by the production of

15 SEAP by vector 5 (or by the production of  $\beta$ -galactosidase by the exchange cassette of Fig. 8B) in response to the indirect or direct regulation of the primary genetic loci 3.

The isolation and identification of trapped regulatory elements as described herein allows the identification of genes operably linked to the trapped regulatory elements and thus the identification of genes whose transcription is increased or decreased by a

20 stimulatory agent of interest. The regulation of these genes can be compared under different environmental conditions and in different cell lines (*e.g.*, cells from different tissues, different organisms, or different disease animal models) to determine whether the genes are regulated the same way in various cell types and to determine whether the regulation of the genes is altered in the presence of certain environmental factors or

25 disease states. The selected cells may be further characterized to determine what proteins affect the transcription of the genes (as described in Example 10) or to determine the role of the encoded proteins *in vivo*, such as the role of wild-type or mutant forms of the encoded protein in inhibiting, causing, or enhancing a disease state.

The selected cells may also be classified based on the characteristics of the

30 trapped regulatory elements or trapped genes. For example, cells containing trapped nucleic acids associated with the expression of proteins in a common class of proteins

(e.g., kinases, phosphatases, proteins in the same signal transduction pathway, or proteins associated with the same disease state) may be classified into the same group. A group of cells may be contacted with a candidate compound such as a potential drug product to compare the effect of the candidate compound on each cell, thereby determining whether the affect of the candidate compound is specific for certain trapped nucleic acids or has a general effect on multiple trapped nucleic acids. As illustrated in Fig. 10, the cells can be used to determine whether different ligands act through separate or overlapping pathways.

The cells of the invention are also useful in the identification and validation of new genetic targets for the treatment or prevention of diseases. For example, the cells can be used to determine whether activation or inhibition of a trapped regulatory element or gene of interest modulates a pathway associated with a disease state. These cells can be used in screening assays to identify new drug products or lead compounds for drug development. Cells containing an inserted reporter gene can be used to identify regulatory elements or promoters that are responsive to a pharmaceutically active compound, such as TNF- $\alpha$ . Cell lines may be selected that are responsive to only TNF- $\alpha$  or are also responsive to other pro-inflammatory cytokines. For example, cell clones which respond to both pro-inflammatory cytokines, TNF- $\alpha$  and IL-1 $\beta$ , can be selected by treating TNF- $\alpha$  cell lines with IL-1 $\beta$  in the presence of the positive selection drug. Thus, by establishing a broad library of cell lines, each incorporating a reporter gene at regulated genetic sites and exhibiting a standardized read-out mechanism, a platform can be assembled with the capability to readily test for efficacy and side-effects of compounds targeting regulatory pathways.

Therapeutic agents based on the identified gene (e.g., antisense molecules, gene activators, or gene inhibitors) can then be appropriately devised. For instance, the gene can be used in gene therapy applications when formulated into appropriate vectors tolerated by the patient in a medical therapeutic delivery vehicle. Alternatively, the gene or its regulatory elements, such as promoters and enhancers, can be used as drug targets to identify potential therapeutic candidates from libraries of compounds.

The following examples are illustrative of certain embodiments of the invention, and are intended to further describe the present invention, without limiting it thereby.

Various modifications can be made to these embodiments without departing from the spirit or scope of the invention.

#### EXAMPLE 1

##### Preparation of a Nucleic Acid Construct

5

The retroviral vector containing the insert shown in Fig 1 is prepared in 5 steps. These steps involve the transfer of cDNA fragments coding for the SEAP and the tTA into expression vectors containing IRES, and then the subsequent merger and transfer of these two constructs into a retroviral vector. These steps are as follows:

- 10 Step 1: The SmaI – XbaI fragment from the pSEAP-2 vector (Clontech) is isolated and inserted by ligation into the SmaI-XbaI sites of the vector pIRES (Clontech).
- Step 2: The EcoRI-BamHI fragment from the pTet-on plasmid (Clontech) is inserted by blunt end ligation into the SmaI site of the vector pIRES.
- Step 3: The EcoRI-XbaI fragment from the vector constructed in step 2 is transferred by
- 15 bluntend ligation into the SmaI site of the vector PBSKS (Stratagene).
- Step 4: The EcoRI-EcoRI fragment of the vector constructed in step 3 is transferred into the EcoRI site of the vector constructed in step 1.
- Step 5: The ClaI-ClaI fragment resultant from step 4 is transferred into the ClaI site of the retroviral vector pSIR (Clontech).

20

#### EXAMPLE 2

##### Construction of a Vector Containing the Nucleic Acid Construct of Example 1.

- This vector is constructed in two steps that include replacement of the neomycin
- 25 resistance gene in the “SATEO” construct (U.S. Patent No. 5,922,601) by the Zeocin resistance gene, and its subsequent transfer to the retroviral vector described in Example 1. These steps are:

- Step 1: Isolation of Zeo cDNA EagI –EcoRI fragment from the pEM7/Zeo vector (Invitrogen) and ligation into the Eag I–EcoRI sites of SATEO.
- 30 Step 2: Isolation of XhoI – BamHI fragment from the construct made in step 1, and its ligation into the XhoI – BamHI sites of the vector described in Example 1.

Retrovirus is produced by transfection into the helper 293 packaging cell line as described in the Clontech Manual for the pSIR vector. Retroviral titer is established by measuring the amount of SEAP activity in infected 3t3 fibroblast cells.

5

### EXAMPLE 3

#### Construction of an Alternative Vector

This vector is constructed in three steps involving the initial deletion of the neomycin resistance gene from the "SATEO" construct (U.S. Patent No. 5,922,601), and the transfer of the resultant insert in combination with the insert made in step 4 of Example 1 into the pSIR vector. The steps are:

10

Step 1: Removal of EagI – SalI insert by digestion "SATEO" construct (U.S. Patent No. 5,922,601), and followed by bluntend ligation.

15

Step 2: Transferring the ClaI – ClaI fragment from the construct made in step 4 of Example 1 to the EcoRI site of the vector pSIR vector.

Step 3: Isolation of XhoI – BamHI fragment from construct made in step 1, and its ligation into the XhoI – BamHI sites of the vector described step 2.

Retrovirus is produced and tittered as described in Example 2.

20

### EXAMPLE 4

#### Preparation of Nucleic Acid Construct for Identifying Genes

This construct is synthesized in 5 steps. The first step involves the synthesis and transfer of an IRES-SD fragment and its placement down stream in a neomycin resistance gene. The entire insert is then transferred into the pTRE vector. The pTRE vector and the insert are then transferred into a retroviral vector. The steps are involved are as follows:

25

Step 1: Two complementary oligonucleotides containing the SD site flanked by restriction sites of XbaI-NotI are synthesized: 5'-aatctagaaggtaaggcgccgcaa-3' (SEQ ID

30

NO.: 1) and 5'-ttgcggccgccttacctctagatt-3' (SEQ ID NO.: 2)

Step 2: Oligonucleotides described in step 1 are annealed and cut by restriction enzymes before being ligated into the XbaI-NotI site in the pIRES vector (Clontech).

Step 3: The neomycin gene from the psv2neo construct (Stratagene) is ligated by bluntend into the MluI site of the vector constructed in step 2.

- 5 Step 4: The EcoRI – BamHI fragment of the vector from step 3 is isolated and ligated into the EcoRI-BamHI site of the pTRE vector (Clontech).

Step 5: The XhoI-NotI fragment from the construct synthesized in step 4 is transferred into the XhoI-BamHI site of the pSIR retroviral vector (Clontech).

10

## EXAMPLE 5

### Preparation of Mast Cell Line library

Mast cells are known to play a central role in inflammatory diseases such as asthma. Cytokines, such as Stem cell factor (SCF) and IL-3, are known to be critical for the proliferative and activation response of mast cells. In vivo, these cytokines induce not only the accumulation of mast cells in airways, but also prime the cells and enhance their hyper-responsiveness. The identification of regulatory factors that can modulate mast cell responses by such cytokines are prime targets for inhibitory drugs. Furthermore, identification of regulatory factors that are involved in the regulation of more than one cytokine in mast cells is likely to represent a critical convergent point of different important pathways. Generation and identification of a mast cell line incorporating such regulatory factors would therefore be highly useful for both high throughput screen for inhibitors and as a means for gene discovery.

#### Experimental Procedure

- 25 The human mast cell line HMC-1 is an established cell line that manifests proliferative and activation responses to various cytokines including IL-3 and SCF. Treatment of cells with cytokines can either up or down regulate genes. In this experiment, cell lines are established containing genes that are up-regulated by IL-3 and SCF. (HMC- I cells are normally maintained in culture medium without additional growth factors).
- 30

To trap IL-3 responsive regulatory regions of genes, HMC-1 cells are cultured in medium without growth factor supplement overnight for 12 hours. HMC-1 cells are then cultured in IL-3 containing medium for 6 hours. After this, the cells are infected with a retrovirus carrying the induction gene trap vector described in Example 2 by culturing  
 5 cells in viral-containing medium for 12 hours. Infected cells are washed once and redistributed into 96 well culture plates at cell numbers of 5000-10,000 per 100ul per well. Selection is initiated with zeocin-containing medium. After three days, surviving cells are collected. These cells represent 1) house-keeping genes or 2) genes activated by IL-3 resulting in the promoters driving production of reporter ZEO, and reporter gene  
 10 transcripts and protein. Reporter assays are performed to demonstrate and confirm the specific expression in the surviving clones.

Selection of the IL-3 responsive genes demonstrates the reversibility of IL-3 induction by switching the culture medium to IL-3-minus medium supplemented with gancyclovir. Housekeeping genes that continue to be active are selected against by the  
 15 expression of thymidine kinase resulting in the elimination of these clones. Surviving clones represent IL-3 responsive genes. To confirm this, a reporter assay are repeated 12 hours after IL-3 deprivation. Clones that are reporter negative are identified.

A similar experiment as described above is carried out to establish cell lines that are SCF responsive. A similar experiment is carried out except that IL-3 is used in place  
 20 of SCF.

To confirm the factor-responsiveness of the isolated clones, reporter assays are repeated for each clone before and after induction. The results are further strengthened with titration curves to quantify dose response.

Each IL-3-responsive cell line is tested with SCF to identify cell lines that will  
 25 respond to both cytokines. Similarly, SCF-responsive cell lines are tested with IL-3 .

In the final step, the identity of the gene for each clone is established. Primers have been synthesized that are specific for use in a 5' RACE with the vectors of this invention to allow cloning and sequencing of the trapped gene. From this information, clones that are responsive to 2 or more factors will be identified.

30

## EXAMPLE 6

### Preparation of Endothelial Cell Line library

Inhibition of angiogenesis is a potent approach to eliminate cancerous tissues. Currently, an increasing number of “anti-angiogenic” molecules have been isolated and are in clinical trials. However, their effect on human cancers has not been established. It is also not known whether a single angiogenesis inhibitor will suffice to maintain the persistent suppression of cancer growth. It is likely that eventually, a combination of therapeutic inhibitors may be necessary. This is not surprising since extensive experimental data has demonstrated that several factors have the capacity to induce proliferation of endothelial cells and promote the genesis of new blood vessels. These factors originated both from the cancer cells and the surrounding stromal components. Gene products modulating these events represent prime targets for inhibitory drugs and small molecules. Similarly, the group of genes that are responsive to more than one factor likely represent critical convergent points of different important pathways. Such cell lines would therefore represent a highly useful tool in a high throughput screen for inhibitors.

#### Experimental procedure

Several well studied factors are known to induce endothelial cell growth. Clones responsive to VEGF, TGF- $\beta$  and FGF-2 are established. The human endothelial line ECV304 or HMEC1 that has been extensively used in other experiments is utilized. Endothelial cells are plated out in 96 well plates at sub-confluent cell density. Cells are stimulated with VEGF-containing medium for 6 hours followed by infection with medium containing retroviral -vectors as described above. 12 hours after initiation of infection, the culture medium is replaced with zeocin-containing medium to select for trapped active genes. As described above, after three to four days of selection, surviving cells represent 1) house-keeping genes or 2) genes induced by VEGF. Reporter assays are performed to demonstrate and confirm the specific expression of the reporter gene in the surviving clones.

To select for VEGF-responsive regulatory regions of genes, the reversibility of VEGF induction is demonstrated by switching the culture medium to medium without VEGF, and supplemented with gancyclovir. Reporter assays are performed 12 hours after

VEGF deprivation. Clones that become reporter-negative are identified.

Reporter-positive clones representing housekeeping genes that continued to be actively transcribed are selected against by the expression of thymidine kinase resulting in the elimination of these clones. Surviving clones after 3-4 days represent VEGF-responsive genes.

To establish cell lines that are TGF- $\beta$  or FGF-2 responsive, a similar experimental sequence as described above for VEGF is carried out, except that TGF- $\beta$  or FGF-2 is used in place of VEGF.

To confirm the specific factor-responsive characteristic of the isolated clones, reporter assays are repeated for each clone before and after induction. Using primers as discussed in previous section in 5' RACE analysis, the identity of the trapped genes is established. Clones that represent genes in endothelial cells responsive to all three stimulants are thus identified.

## EXAMPLE 7

### Selection of a Gene which up regulates or down regulates the selected Loci

Selected clones from Examples 5 and 6 are cultured in a growth medium until 80% confluency is reached. These cells are then infected with a retrovirus carrying the gene trap vector described in Example 4 by culturing cells in viral containing medium for 12 hours. Infected cells are washed once and redistributed into 96 well culture plates at cell numbers of 5000-10,000 per 100ul per well. Selection is initiated with G418-containing medium. After three days, surviving cells are collected. These cells represent cells in which the viral vector has been successfully integrated. The clones from Example 5 are placed in growth containing medium containing Zeocin and G418. This allows for the selection of cells with active genomic loci, in addition to an integrated gene trap vector which confers resistance to G418. Reporter assays are performed to demonstrate and confirm the specific expression in the surviving clones.

The clones from Example 6 are selected based on testing for ligand independent SEAP activity. For example, these cells may be incubated in the absence of the stimulatory agent to identify trapped genes that encode proteins which modulate the

activity of the regulatory element operably linked to the SEAP coding sequence in a ligand independent manner.

The cells may also be incubated in the presence of the stimulatory agent to identify trapped genes that encode proteins which modulate the activity of the regulatory element in a ligand dependent manner. In particular, the encoded proteins that are more active in the presence of the stimulatory agent produce a greater effect on the level of SEAP activity in the presence of the stimulatory agent. These encoded proteins may be directly activated by the stimulatory agent or may be activated by another protein which is directly or indirectly activated by the stimulatory agent. Alternatively, the stimulatory agent may inhibit another protein that would otherwise inhibit the protein encoded by the trapped gene. The encoded proteins that are less active in the presence of the stimulatory agent produce a smaller effect on the level of SEAP activity in the presence of the stimulatory agent. These encoded proteins may be directly or indirectly inhibited by the stimulatory agent.

Validity of the model is tested by looking for clones that demonstrate SEAP production in a tetracycline-dependent manner. For example, trapped genes that encode proteins which activate a regulatory element of interest enhance the production of SEAP in the presence of tetracycline. Conversely, trapped genes that encode proteins which inactivate a regulatory element of interest inhibit the production of SEAP in the presence of tetracycline. Using primers corresponding to sequences upstream of the SD site in the gene trap vector in 3' RACE analysis, the identity of the trapped genes is established.

## EXAMPLE 8

### Screening for Inhibitors and Antagonists

Selected cell clones from Examples 6 and 7 are used directly to screen a natural products library (EXALPHA) for inhibitors and activators of SCF and VEGF activity. For identification of inhibitors to the SCF mediated signaling in the HMC-1 clones, these cells were cultured and plated equally into eleven 96 well plates. For identification of inhibitors and activators, a 1 nM aliquot from each well of the natural products library

were transferred to each well of cultured cells in the presence of SCF. The amount of SEAP activity is measured and compared in well-to-well manner. For identification of SCF independent activators, this screen is done in the absence of SCF.

- For identification of inhibitors and activators of the VEGF mediated signaling in ECV304 cells, similar techniques as above are used. ECV304 cell clones are harvested and plated in 96 well plates, and the relative amount of SEAP produced is compared in each well in the presence of 1 nM of the natural product fraction and VEGF.

### EXAMPLE 9

#### Identification of Regulatory Elements that are Responsive to a Stimulatory Agent

- For the identification of regulatory elements that are responsive to a stimulatory agent of interest, cells are infected with a retrovirus carrying the induction gene trap vector illustrated in Fig. 8A or Fig. 9A or a similar vector containing one or no LoxP sites, as described in Example 2. The infected cells are then washed once and redistributed into 96 well culture plates.

#### *Selection of Regulatory Elements that are Activated by a Stimulatory Agent*

- To identify regulatory elements (*e.g.*, enhancers or promoters) that are activated by a stimulatory agent of interest, the cells are incubated in the presence of the stimulatory agent of interest and Zeocin, the positive selection drug (Fig. 6). This step results in the isolation of cells in which the construct has stably integrated into the genome under the control of a promoter that may or may not be regulated by the stimulating agent. To eliminate cells in which the construct is expressed under the control of a promoter that is not regulated by the stimulating agent (*e.g.*, a housekeeping gene promoter), the cells are cultured in the absence of the stimulating agent, but in the presence of gancyclovir. This step eliminates cells that express the negative selective marker thymidine kinase in the absence of the stimulatory agent and results in the isolation of desired cells in which the construct has stably integrated into the genome under the control of a promoter (or other regulatory element) that is regulated by the stimulating agent. While not meant to limit the invention in any way, it is noted that the

integration of the construct into the cells is an essentially random event, thus not all of the cells will contain a construct integrated under the control of a endogenous promoter or under the control of an endogenous promoter modulated by the stimulating agent of interest.

5           Another method that may be used to identify regulatory elements that are activated by the stimulatory agent involves first incubating the cells with Zeocin to select cells containing the induction trap vector (Fig. 7). The selected cells are then incubated in the presence of gancyclovir without the stimulatory agent. This step eliminates undesired cells in which the trapped regulatory elements are transcriptionally active in the absence  
10 of the stimulatory agent. The remaining cells are incubated in the presence of the stimulatory agent. The desired cells that are responsive to the stimulatory agent are selected based on the transcription of the reporter gene (*e.g.*, by measuring SEAP production). Thus, the reporter gene from the induction trap vector allows the effect of the stimulatory agent on the regulatory element to be quantitated. This quantitation  
15 allows the effect of different stimulatory agents on the same cell to be compared and allows the effect of one stimulatory agent on different cells to be compared. In desirable embodiments, the effect of a stimulatory agent of interest is at least 2, 5, 8, 10, 20, 50, or 100 fold greater than the effect of another stimulatory agent on the transcription of the reporter gene. In other embodiments, the effect of a stimulatory agent of interest is at  
20 least 2, 5, 8, 10, 20, 50, or 100 fold greater than the effect of the stimulatory agent on a corresponding control cell that lacks the regulatory element of interest or that has regulatory elements with polynucleotide sequences that are less than 60, 40, 30, 20, or 10% identical to the polynucleotide sequence of the regulatory element of interest.

#### 25   *Selection of Regulatory Elements that are Inhibited by a Stimulatory Agent*

For the identification of regulatory elements (*e.g.*, enhancers or promoters) that are inhibited by a stimulatory agent of interest, the cells are incubated in the presence of Zeocin to select cells containing the induction trap vector (Fig. 6). Then, the selected cells are incubated in the presence of both the stimulatory agent and gancyclovir. This  
30 incubation eliminates undesired cells containing trapped regulatory elements that are transcriptionally active in the presence of the stimulatory agent, allowing cells in which

the trapped regulatory elements are inactivated by the stimulatory agent to be selected. The selected cells may be assayed to confirm that the reporter gene is transcribed in the absence of the stimulatory agent, resulting in SEAP production (Fig. 7).

## 5 *Identification of Trapped Genes*

The sequence of the trapped regulatory elements that are upstream of the integrated construct may be determined using standard 5' RACE molecular biology methods, as described in Example 5. Additionally, the coding sequence for the trapped gene that is upstream and/or downstream of the integrated construct may be determined  
10 using standard DNA amplification and sequencing methods.

Alternatively, if an induction trap vector is used that contains a prokaryotic promoter (*e.g.*, a bacterial promoter) operably linked to the positive selection marker, such as the vector illustrated in Fig. 9A, bacterial cells may be used to facilitate the identification of the trapped regulatory elements. In this method, genomic DNA from a  
15 selected eukaryotic cell is isolated and digested with a restriction enzyme that cleaves the integrated construct at one site and cleaves the endogenous, eukaryotic genomic DNA flanking the integrated construct at one or more sites. Or the DNA is digested with a restriction enzyme that does not cleave the integrated construct but cleaves the endogenous, eukaryotic genomic DNA at two or more sites. Alternatively, two restriction  
20 enzymes can be used so that one restriction enzyme cleaves the endogenous DNA and the other restriction enzyme cleaves either another site in the endogenous DNA or cleaves a site in the integrated construct.

For example, for the vector illustrated in Fig. 9A, the *Cla*I restriction enzyme is used to cleave the integrated construct at a single, predetermined site and to cleave the  
25 eukaryotic genomic DNA at one or more cleavage sites. The restriction enzyme-digested DNA fragments are then ligated to a restriction enzyme digested bacterial plasmid. The desired, ligated bacterial plasmids contain an insert with the positive selection marker (*e.g.*, zeocin) from the construct that integrated into the selected eukaryotic cell and contain a region of the eukaryotic genomic DNA flanking the integrated construct. To  
30 select these desired plasmids, the plasmids are used to transform competent bacterial cells, and the transformed bacterial cells are grown on plates containing the selection

agent to which the positive selection marker present within the insert confers resistance (*e.g.*, zeocin). If a bacterial plasmid that also contains an endogenous positive selection marker is used, such as the PBSK vector that contains the ampicillin drug resistance gene, the transformed bacteria can be plated on plates containing both positive selection agents

5 (*e.g.*, ampicillin and zeocin). The selected bacteria contain a region from the eukaryotic genomic DNA that flanked the integrated induction trap vector. The size of this eukaryotic genomic DNA fragment can be calculated based on the size of the insert that was added to the bacterial plasmid (*e.g.*, based on the migration in an agarose gel compared to the migration of standards with known molecular weights). The sequence of

10 this eukaryotic genomic DNA can be readily determined by PCR amplifying and sequencing the insert in the bacterial plasmid using a primer designed to bind a region of the plasmid, such as a primer that binds the prokaryotic promoter upstream from the insert, or using a primer designed to bind a region in the insert. The sequence of the genomic DNA can be compared to known sequences, such as the publicly available

15 sequence of the human genome, to identify the eukaryotic regulatory elements trapped by the induction trap vector and to identify the genes that are operably linked to these regulatory elements.

Similarly, if an induction trap vector is used that contains a yeast promoter operably linked to the positive selection marker, yeast cells may be used to facilitate the

20 identification of the trapped regulatory elements. This method is performed essentially as described above, except that yeast cells are transformed with a plasmid containing a yeast promoter and an insert which includes the positive selection marker and a region of the eukaryotic, genomic DNA that flanked the integrated induction trap vector in the selected eukaryotic cells. The yeast cells containing the desired plasmid are selected using the

25 positive selection agent, and then the insert is PCR amplified and sequenced as described above.

## EXAMPLE 10

### Identification of Genes Encoding Proteins which Modulate Regulatory Elements

As described in Example 7, genes may be identified that encode proteins which modulate the transcriptional activity of the regulatory elements identified using an induction trap vector. In one possible method, a transactivator coding sequence (*e.g.*, teton/off) is added to the region of the induction trap vector that integrated into the genome of the isolated cells from Example 9. Any standard molecular biology technique may be used to add this transactivator coding sequence.

### *Cassette Exchange*

For example, a vector containing an exchange cassette with the transactivator coding sequence and a reporter gene flanked by LoxP sites may be used to replace the region of the induction trap vector of Example 9 that is flanked by LoxP sites. A LoxP site consists of a double-stranded 34 basepair sequence. This sequence contains two 13 basepair inverted repeat sequences that are separated from one another by an 8 basepair spacer region (Hoess *et al.*, Proc. Natl. Acad. Sci. U.S.A. 79:3398-3402, 1982; Sauer, U.S. Patent No. 4,959,317). One strand of the LoxP site has the sequence 5'-ATAACTTCGTATAATGTATGCTATACGAAGTTAT-3' (SEQ ID NO.:3), and the other strand has the sequence 5'-ATAACTTCGTATAGCATACATTATACGAAGTTAT-3' (SEQ ID NO.:4). Alternatively, other lox sites (*e.g.*, Lox 511 sites) or LoxP sites containing nucleotide substitutions that do not prevent recognition by the Cre recombinase may be used (Sauer, Methods: A Companion to Methods in Enzymology 14:381-392, 1998).

This Cre recombinase-mediated cassette exchange may be performed by transfecting the selected cells from Example 9 with the vector illustrated in Fig. 8B that contains the LoxP flanked exchange cassette and with a vector encoding Cre recombinase (see, for example, Fukushige and Sauer, Proc. Natl. Acad. Sci. USA 89:7905-7909, 1992; Feng *et al.*, J. Mol. Biol. 292:779-785, 1999; U.S. Patent No. 4,959,317; Proc. Natl. Acad. Sci. U.S.A. 85:5166-5170, 1988). Alternatively, the selected cells may be transfected with a vector that contains both the LoxP flanked exchange cassette and a Cre recombinase coding sequence. The cells in which Cre-mediated recombination has taken place may be selected based on the expression of the reporter gene from the exchange cassette (*e.g.*,  $\beta$ -galactosidase) and based on Zeocin sensitivity. Expression of the

transactivator polypeptide may also be confirmed by western blotting. The above method may also be used if one or both of the vectors contain only one LoxP site.

Alternatively, the cassette exchange may be performed using recombinase signal sequences and a recombinase from any other site-specific recombinase system. For example, the flp recombinase (Schwartz *et al.*, J. Molec. Biol. 205:647-658, 1989; Parsons *et al.*, J. Biol. Chem. 265:4527-4533, 1990; Golic *et al.*, Cell 59:499-509, 1989; Amin *et al.*, J. Molec. Biol. 214:55-72, 1990); the site-specific recombination system of the *E. coli* bacteriophage  $\lambda$  (Weisberg *et al.*, In: Lambda II, (Hendrix *et al.*, Eds.), Cold Spring Harbor Press, Cold Spring Harbor, N.Y., pp. 211-250 (1983), TpnI and the  $\beta$ -lactamase transposons (Levesque, J. Bacteriol. 172:3745-3757, 1990); the Tn3 resolvase (Flanagan *et al.*, J. Molec. Biol. 206:295-304, 1989; Stark *et al.*, Cell 58:779-790, 1989); the yeast recombinases (Matsuzaki *et al.*, J. Bacteriol. 172:610-618, 1990); the *B. subtilis* SpoIVC recombinase (Sato *et al.*, J. Bacteriol. 172:1092-1098, 1990); the Hin recombinase (Glasgow *et al.*, J. Biol. Chem. 264:10072-10082, 1989); immunoglobulin recombinases (Malynn *et al.*, Cell 54:453-460, 1988); or the Cin recombinase (Haftner *et al.*, EMBO J. 7:3991-3996, 1988; Hubner *et al.*, J. Molec. Biol. 205:493-500, 1989) can be used. These alternative systems are also discussed by Echols (J. Biol. Chem. 265:14697-14700, 1990), de Villartay (Nature 335:170-174, 1988), Craig (Ann. Rev. Genet. 22:77-105, 1988), Poyart-Salmeron *et al.* (EMBO J. 8:2425-2433, 1989), Hunger-Bertling *et al.* (Molec. Cell. Biochem. 92:107-116, 1990), and Cregg (Molec. Gen. Genet. 219:320-323, 1989).

The region of the induction trap vector that is replaced by the exchange cassette includes an IRES site; thus, replacing this region with the exchange cassette, rather than adding the exchange cassette downstream or upstream of this region, results in the elimination of this IRES site. Because multiple IRES sites near the reporter gene may decrease the transcription of the reporter gene, eliminating this IRES site may result in greater reporter gene expression than the corresponding level of reporter gene expression if this IRES site is maintained. The reporter gene in the exchange cassette (*e.g.*,  $\beta$ -galactosidase) may be the same or may be different from that of the induction trap vector. Alternatively, either the induction trap vector or the exchange cassette may contain a reporter gene and the other one may lack a reporter gene. For example, if the exchange

cassette does not contain a reporter gene, the integration of the exchange cassette into the genome of the cells may be determined by northern or western blotting for the encoded transactivator mRNA or protein.

The exchange cassette or the induction trap vector may optionally contain a  
 5 prokaryotic or yeast promoter operably linked to a reporter gene or a positive selection  
 marker to allow a region from the integrated construct and a region of the flanking  
 eukaryotic, genomic DNA to be transferred to a bacterial or yeast plasmid, as described in  
 Example 9. The bacterial or yeast plasmid can be easily produced in large quantities by  
 the growth of bacteria or yeast transformed with the plasmid, and then PCR-amplified and  
 10 sequenced to identify the trapped regulatory elements.

#### *Introduction of Gene Trap Vector*

In addition to undergoing this cassette exchange, the cells are also transfected with  
 a gene trap vector that includes a tetracycline responsive element operably linked to a  
 15 minimal promoter (*e.g.*, TRE<sub>pminCMV</sub>), a positive selective marker (*e.g.*, Neo), an IRES  
 sequence, and a splice donor. An exemplary construct is illustrated in Fig. 4. The gene  
 trap vector may optionally contain a prokaryotic or yeast promoter operably linked to the  
 positive selection marker. Transfected cells containing this construct may be selected  
 using the positive selection drug to which the construct confers resistance. This positive  
 20 selection marker may be the same or may be different from the positive selection marker  
 in the induction trap vector encoding the transactivator.

Alternatively, a gene trap vector without a positive selection marker may be used.  
 For example, cells containing a regulatory element that is activated by a stimulatory agent  
 may be incubated in the absence of the stimulatory agent. Under these conditions, there  
 25 is little or no expression of the positive selection marker in the induction trap vector  
 because the stimulatory agent is not present to activate the endogenous regulatory element  
 of interest that controls the expression of the positive selection marker. The gene trap  
 vector is then inserted into the cells. In some or all of the cells, the TRE<sub>pminCMV</sub> promoter  
 from this vector integrates into the genome of the cells such that it is operably linked to an  
 30 endogenous gene encoding a protein that activates the regulatory element of interest. The  
 residual promoter activity of the TRE<sub>pminCMV</sub> promoter in the absence of the stimulatory

agent activates the transcription of the trapped gene, and then the encoded protein activates the expression of the positive selection marker. Thus, cells containing the gene trap vector may be selected based on the increased expression of the positive selection marker in the induction trap vector.

5           Similarly, cells containing a regulatory element that is inactivated by a stimulatory agent may be incubated in the presence of the stimulatory agent. Under these conditions, there is little or no expression of the positive selection marker in the induction trap vector because the stimulatory agent inhibits the endogenous regulatory element controlling the expression of the positive selection marker. If the TRE<sub>pminCMV</sub> promoter from the gene  
10   trap vector integrates into the genome of the cells upstream of an endogenous gene encoding a protein that activates the regulatory element of interest, the encoded protein activates the expression of the positive selection marker, allowing cells containing the gene trap vector to be selected based on the increased expression of the positive selection marker.

15

#### *Selection of Genes Encoding Proteins that Activate Regulatory Elements of Interest*

To identify genes that activate the regulatory elements discovered in Example 9, the cells containing the exchange cassette and the gene trap vector are cultured in the presence of tetracycline, which forms a complex with the protein encoded by the teton/off  
20   nucleic acid. This complex activates expression of genes downstream of minimal promoters including tetracycline responsive elements. Thus, if the gene trap vector has integrated upstream of a gene encoding a protein that activates the regulatory element of interest, the encoded protein increases the level of transcription of the reporter gene (e.g.,  $\beta$ -galactosidase) that is downstream of the regulatory element of interest. Culturing these  
25   cells in the presence of tetracycline leads to greater expression of the reporter gene than the corresponding level in the absence of tetracycline. These desired cells are selected based on their increased level of reporter gene expression or activity.

#### *Selection of Genes Encoding Proteins that Inhibit Regulatory Elements of Interest*

30           For the identification of genes encoding proteins that inactivate the regulatory elements discovered in Example 9, the cells are also cultured in the presence of

tetracycline. Cells in which the gene trap vector has integrated upstream of a gene encoding a protein that inhibits the regulatory element of interest have lower levels of reporter gene expression in the presence of tetracycline than in the absence of tetracycline. Thus, these desired cells may be selected based on the inhibition of reporter gene expression or activity.

### *Identification of Trapped Genes*

The sequence of the trapped genes that are downstream of the integrated construct may be determined using standard DNA amplification and sequencing methods.

- Alternatively, if the gene trap vector contains a prokaryotic or yeast promoter operably linked to a positive selection marker, bacterial or yeast cells may be used to facilitate the identification of the trapped genes as described in Example 9.

## EXAMPLE 11

### Selection of Cell Lines Responsive to Pro-Inflammatory Ligands

Cell lines were generated that are responsive to ligands in pro-inflammatory pathways involved in rheumatoid arthritis (RA), which is an auto-immune disease associated with recurrent and progressive pain and inflammation of joints. NSAID agents are commonly used to reduce the pain and signs of inflammation in rheumatoid arthritis patients. Cells involved in rheumatoid arthritis include fibroblasts and CD4<sup>+</sup> T cells. Members of the cytokine and chemokine signaling pathway in rheumatoid arthritis include TNF $\alpha$ , IL-6, IL-1 $\beta$ , and SDF-1.

- As illustrated in Fig. 11, the methods described herein were used to isolate EL4 or NIH3T3 fibroblast cells activated by TNF $\alpha$  or IL-1 $\beta$ . To confirm that the reporter gene (SEAP) that integrated into the genome of these cells was integrated under the control of a regulatory element responsive to TNF $\alpha$  or IL-1 $\beta$ , the selected cells were exposed to TNF $\alpha$  or IL-1 $\beta$ , and SEAP activity was measured (Fig. 12A). As expected, SEAP activity was induced by IL-1 $\beta$  in NIH3T3 cells selected for their responsiveness to IL-1 $\beta$  and by TNF $\alpha$  in EL4 cells selected for their responsiveness to TNF $\alpha$ . SEAP activity was also induced by SDF-1 in some of the cells selected for their responsiveness to TNF $\alpha$ . As

illustrated in Fig. 12B, a probe to the TK/Zeo selection markers in the integrated construct was used in standard southern blot analysis to confirm the integration of the construct in some of the selected NIH3T3 cell lines.

#### The NIH3T3

5 cells selected for their responsiveness to IL-1 $\beta$  were tested to determine whether they were also responsive to other pro-inflammatory molecules. As illustrated in Figs. 13A and 13B, seven clones had the highest level of responsiveness to IL-1 $\beta$ , based on SEAP activity. The clones had varying levels of responsiveness to TNF $\alpha$  and IL-6. The clones that were responsive to all three ligands demonstrate that some of the pathways activated  
10 by IL-1 $\beta$ , TNF $\alpha$ , and IL-6 overlap. The clones, such as D5, that were responsive to IL-1 $\beta$  but had negligible response to TNF $\alpha$  and IL-6 demonstrate that there are IL-1 $\beta$  specific pathways that are independent of TNF $\alpha$  and IL-6.

Similarly, EL-4 clones selected for their responsiveness to TNF $\alpha$  were tested to determine whether they were responsive to other ligands. Two clones were also  
15 responsive to IL-1, PMA, and IL-10; in order of decreasing responsiveness (Fig. 14A). EL-4 clones selected for their responsiveness to IL-10 were also responsive to TNF $\alpha$  and IL-1 (Fig. 14B). These results indicate that there are overlapping pathways involving TNF $\alpha$ , IL-1, PMA, and IL-10. EL4 cells responsive to both TNF $\alpha$  and SDF-1 were selected by treating TNF- $\alpha$  cell lines with SDF-1 in the presence of the positive selection  
20 drug (Fig. 14C).

## EXAMPLE 12

### Demonstration of a Link between Cox-2 Activity and a Pro-Inflammatory Signaling Pathway

25

As illustrated in Fig. 15, the specific Cox-2 inhibitor, celecoxib, was shown to inhibit the effect of IL-1 $\beta$  on SEAP reporter gene activity in selected NIH3T3 cells in a concentration dependent manner. The IC<sub>50</sub> value of the inhibition of SEAP activity by celecoxib was approximately 0.2  $\mu$ M. In contrast, celecoxib was ineffective at inhibiting  
30 the effect of TNF $\alpha$  on SEAP reporter gene activity in selected EL-4 cells. In some

clones, TNF $\alpha$  increased SEAP reporter gene (Fig. 16). The clones not affected by the celecoxib include PD-5, PA-6, PA-5, and PB-5. The clones affected by celecoxib include PD-6 and C-5. Thus, within a targeted cell type, some clones are affected by the Cox-2 inhibitor and some clones are not affected.

5 As illustrated in Figs. 17A and 17B, different selected EL-4 clones had different levels of response to various ligands and ligand combinations. These results indicate that candidate drug products may have effects in multiple pathways. In some cases, the effect of a candidate drug in one or more pathways leads to adverse side-effects when the drug is administered to mammals (*e.g.*, humans). Thus, candidate drugs that are identified as  
10 activating a pathway associated with adverse effects, such as toxic effects or the promotion of a disease state, are desirably eliminated from further drug development. Similarly, candidate compounds identified as inhibiting a pathway associated with beneficial effects (*e.g.*, the reduction of adverse effects, the prevention of a disease state, or the inhibition of the progression of a disease state) are desirably eliminated from  
15 further drug development.

### EXAMPLE 13

#### Use of Selected Cells to Measure Drug Efficacy

20 The NIH3T3 cells selected for their responsiveness to IL-1 $\beta$  were also tested to measure the efficacy of the MEK inhibitor U0126 and cyclosporin A. As illustrated in Fig. 18A, U0126 inhibited the effect of IL-1 $\beta$  on SEAP reporter gene activity in selected NIH3T3 cells in a concentration dependent manner. The IC<sub>50</sub> value of the inhibition of SEAP activity by MAP kinase inhibitors U0126 and PD98059 was approximately 1.0  
25  $\mu$ M. Cyclosporin A had a much smaller effect on SEAP activity (Fig. 18B). Thus, these selected cells are useful for measuring the activity of candidate drug products in cell-based assays.

## EXAMPLE 14

## Development of bioassays for mutagenic agents

The genetic integrity of DNA is constantly being challenged by an array of DNA damaging agents, which can be either endogeneous or exogeneous in origin. Cellular repair systems are present to counteract potentially mutagenic or cytotoxic consequences from the DNA damage. Base damage is repaired either directly, through dealkylation, or via complex and coordinated pathways involving multiple proteins. These latter systems include mismatch repair (MMR), base excision repair (BER) and nucleotide excision repair. In addition, cellular regulatory pathways are activated by damaged DNA and can serve as a reporter system for the presence of mutagenic agents.

Development of cell lines that report activity of regulatory pathways activated upon exposure of cells to DNA damaging agents such as alkylating agents enables the development of an early screen against such agents and can be used to identify compounds that cause damage to DNA.

For development of such assays a library of NIH3t3 fibroblast cell lines were generated that are responsive to the presence of the DNA-alkylating agent methyl methanesulphonate (MMS). These cells were exposed to MMS (0.2nM) in the presence of the virus made by the viral construct as described in example 2, and the positive selection drug phelomycin. Cells surviving this selection were rested for 2 days before treatment with the negative selection marker Gancyclovir in the absence of MMS. Cell clones that demonstrated inducible SEAP reporter response upon treatment with MMS were isolated. The SEAP reporter response to other DNA mutagens were tested and clones which showed consistent response to such agents were chosen.

Genes down regulated by the presence of DNA mutagens can also be identified by reversing the order of treatment with the positive and negative selection drugs.

*Identification of Trapped Genes*

The sequence of the trapped regulatory elements upstream of the integrated construct may be determined using standard 5' RACE molecular biology methods, as described in Example 5. Additionally, the coding sequence for the trapped gene (mRNA)

that is upstream and/or downstream of the integrated construct may be determined using standard cDNA amplification and sequencing methods. Identification of mRNA's regulated by DNA mutagens will enable the isolation of the regulatory sequences (promoters) of these genes. Constructs utilizing promoters of these regulated genes can be made to drive expression of reporter genes. These constructs can be transfected into cells and the cells used as reporter cells for DNA damaging agents. Also, other techniques such as differential display, PCR select (Clontech) and DNA chip can be utilized to identify genes regulated by DNA mutagens. Monitoring the expression of these genes or their products, in addition to the activity of their promoters, can be used either directly or indirectly as markers for the presence of DNA damaging agents in cells.

Alternatively, if an induction trap vector is used that contains a prokaryotic promoter (*e.g.*, a bacterial promoter) operably linked to the positive selection marker, such as the vector illustrated in Fig. 9A, bacterial cells may be used to facilitate the identification of the trapped regulatory elements. In this method, genomic DNA from a selected eukaryotic cell is isolated and digested with a restriction enzyme that cleaves the integrated construct at one site and cleaves the endogenous, eukaryotic genomic DNA flanking the integrated construct at one or more sites. Or the DNA is digested with a restriction enzyme that does not cleave the integrated construct but cleaves the endogenous, eukaryotic genomic DNA at two or more sites. Alternatively, two restriction enzymes can be used so that one restriction enzyme cleaves the endogenous DNA and the other restriction enzyme cleaves either another site in the endogenous DNA or cleaves a site in the integrated construct. These constructs would contain the regulatory sequence (promoter) of the gene regulated and these DNA can be sequenced and identified. Reporter constructs can be engineered that contain these sequences and they can be transfected into eukaryotic cells and the cells can be used as assays for the presence of DNA mutagens.

## OTHER EMBODIMENTS

Each of the foregoing patents, patent applications and references that are recited in this application are herein incorporated in their entirety by reference. Having described  
5 the presently preferred embodiments, and in accordance with the present invention, it is believed that other modifications, variations and changes will be suggested to those skilled in the art in view of the teachings set forth herein. It is, therefore, to be understood that all such variations, modifications, and changes are believed to fall within the scope of the present invention as defined by the appended claims.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
22